



Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2332533 A1 2001/08/18

(21) **2 332 533**

(12) **DEMANDE DE BREVET CANADIEN  
CANADIAN PATENT APPLICATION**

(13) **A1**

(22) Date de dépôt/Filing Date: 2001/02/16

(41) Mise à la disp. pub./Open to Public Insp.: 2001/08/18

(30) Priorité/Priority: 2000/02/18 (60/184,152) US

(51) Cl. Int. / Int. Cl. : C 12N 15/57, A61K 38/48, C 12N 9/64,  
C 12N 5/00, C 07K 16/40, C 12N 5/10, C 12Q 1/00,  
C 12P 21/00

(71) Demandeur/Applicant:  
F. HOFFMANN-LA ROCHE AG, CH

(72) Inventeurs/Inventors:  
ZUO, FENG RONG, US,  
KLONOWSKI, PAUL, US,  
HELLER, RENU ANAND, US

(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : NOUVELLES METALLOPROTEASES COMPORTANT DES DOMAINES DE NATURE THROMBOSPONDINE  
ET COMPOSITIONS D'ACIDES NUCLEIQUES CODANT CES COMPOSES

(54) Title: NOVEL METALLOPROTEASES HAVING THROMBOSPONDIN DOMAINS AND NUCLEIC ACID  
COMPOSITIONS ENCODING THE SAME

(57) **Abrégé/Abstract.**

Novel metalloproteases having thrombospondin domain(s) (MPTS proteins) and polypeptides related thereto, as well as nucleic acid compositions encoding the same, are provided. The subject polypeptide and nucleic acid compositions find use in a variety of applications, including diagnostic applications, therapeutic agent screening applications, as well as therapeutic applications for a variety of different conditions. Also provided are methods of treating disease conditions associated with aggrecanase activity, e.g. conditions characterized by the presence of aggrecan cleavage products, such as rheumatoid- and osteo-arthritis.





Abstract

Novel metalloproteases having thrombospondin domain(s) (MPTS proteins) and polypeptides related thereto, as well as nucleic acid compositions encoding the same, are provided. The subject polypeptide and nucleic acid compositions find use in a variety of applications, including diagnostic applications, therapeutic agent screening applications, as well as therapeutic applications for a variety of different conditions. Also provided are methods of treating disease conditions associated with aggrecanase activity, e.g. conditions characterized by the presence of aggrecan cleavage products, such as rheumatoid- and osteo-arthritis.



The field of the invention is proteases, particularly metalloproteases with thrombospondin domains.

Cartilage matrix structure as dry weight of the tissue is made up of 70% collagen  
5 and 20-30% proteoglycans. The proteoglycan component confers mechanical flexibility  
to load bearing tissues and imparts viscoelastic properties to cartilage. Its loss leads to  
rapid structural damage as is seen most frequently in arthritic joint diseases and joint  
injury.

10 Aggrecan is a major cartilage proteoglycan. Aggrecan is a large protein of 210  
kDa and has three globular domains: G1, G2, and G3. The G1 and G2 domains of the  
protein are closer to the amino terminus of the protein and their intervening  
interglobular domain has sites that are proteolytically sensitive. The region between G2  
and G3 is heavily glycosylated and connected to oligosaccharides and  
15 glycosaminoglycans (GAGs) to form the mature proteoglycan. In arthritic cartilage,  
core protein fragments of 55 kDa are observed and believed to be the result of cleavage  
of the core protein in the G1 and G2 interglobular domain between asparagine 341 and  
phenylalanine 342. This cleavage can be made by many matrix metalloproteinases e.g.  
MMP-1, -2, -3, -7, -8, -9, and -13. In addition, 60 kDa aggrecan fragments with a -  
20 COOH terminus of glutamic acid are also identified and are indicative of a cleavage site  
between glutamic acid 373 and alanine 374. Matrix metalloproteinase are unable to  
cleave at this site. The unique endopeptidase activity responsible for this cleavage has  
been termed "aggrecanase."

25 The G1 domain of the core protein forms a stable ternary complex by binding to  
hyaluronic acid and link proteins in the matrix. Any enzymatic cleavage in this region  
destabilizes the cartilage matrix structure, leads to the loss of the major proteoglycan  
aggrecan and exposes type II collagen to collagenases, causing cartilage loss and the  
consequent development of joint disease. Since a variety of anti-arthritic drugs do not

As such, aggrecanase is considered to be an important drug target for arthritis. Aggrecan fragments released into the synovial fluid are the primary detectable events in the development of rheumatoid- and osteo- arthritis. Search for this protease has been intense. Despite these intense discovery efforts, identification of human aggrecanase has  
 5 remained elusive.

As such, there is much interest in the identification of human aggrecanase, as well as the gene encoding this activity.

10 The following references are directed to this field. U.S. Patents 5,872,209 and 5,427,954, and WO 99/09000; WO 98/55643; WO 98/51665, and WO 97/18207.

Other references include: Abbasdale, "Cloning and characterization of ADAMTS11, an aggrecanase from the ADAMTS family," *J. Biol. Chem.* (Aug. 1999) 274: 23443-50; Arner et al., "Generation and Characterization of Aggrecanase. A  
 15 soluble, cartilage-derived aggrecan-degrading activity," *J Biol Chem* (1999 Mar 5) 274(10):6594-6601; Arner et al., "Cytokine-induced cartilage proteoglycan degradation is mediated by aggrecanase," *Osteoarthritis Cartilage* (1998 May) 6(3):214-28; Illington et al., "An aggrecan-degrading activity associated with chondrocyte  
 20 membranes," *Biochem J* (1998 Nov 15) 336 ( Pt 1):207-12; Buttner et al., "Membrane type 1 matrix metalloproteinase (MT1-MMP) cleaves the recombinant aggrecan substrate rAgg1mut at the 'aggrecanase' and the MMP sites. Characterization of MT1-MMP catabolic activities on the interglobular domain of aggrecan," *Biochem J* (1998 Jul 1) 333 ( Pt 1):159-65; Flannery et al., "Expression of ADAMTS homologues in articular  
 25 cartilage," *Biochem. Biophys. Res. Commun.* (July 1999) 260:318-22; Hurskainen et al., "ADAM-TS5, ADAM-TS6, and ADAM-TS7, Novel members of a New Family of Zinc Metalloproteases," *J. Biol. Chem.* (Sept. 1999) 274: 25555-25563; Hughes et al., "Differential expression of aggrecanase and matrix metalloproteinase activity in  
 chondrocytes isolated from bovine and porcine articular cartilage," *J Biol Chem* (1998  
 30 Nov 13) 273(46):30576-82; Ilic et al., "Characterization of aggrecan retained and lost from the extracellular matrix of articular cartilage. Involvement of carboxyl-terminal processing in the catabolism of aggrecan," *J Biol Chem* (1998 Jul 10) 273(28):17451-8; Kuno et al., "ADAMTS-1 is an active metalloproteinase associated with the extracellular

- matrix," J. Biol. Chem. (June 1999) 274:18821-6; Kuno et al., "ADAMTS-1 protein anchors at the extracellular matrix through the thrombospondin type I motifs and its spacing region," J. Biol. Chem. (May 1998) 273:13912-7; Kuno et al., "The exon/intron organization and chromosomal mapping of the mouse ADAMTS-1 gene encoding an
- 5 ADAM family protein with TSP motifs," Genomics (Dec. 1997) 46:466-71; Kuno et al., "Molecular cloning of a gene encoding a new type of metalloproteinase-disintegrin family protein with thrombospondin motifs as an inflammation associated gene," J. Biol. Chem. (Jan. 1997) 272: 556-62; Sandy et al., "Chondrocyte-mediated catabolism of aggrecan: aggrecanase-dependent cleavage induced by interleukin-1 or retinoic acid can
- 10 be inhibited by glucosamine," Biochem J (1998 Oct 1) 335 ( Pt 1):59-66; Tang & Hong, "ADAMTS: a novel family of proteases with ADAM protease domain and thrombospondin 1 repeats," FEBS Lett. (Feb. 1999) 445:223-5; Tortorella et al., Purification and cloning of aggrecanase-1: a member of the ADAMTS family of proteins," Science (June 1999) 284:1664-6; Vankemmelbeke et al., "Coincubation of
- 15 bovine synovial or capsular tissue with cartilage generates a soluble 'Aggrecanase' activity," Biochem Biophys Res Commun (1999 Feb 24) 255(3):686-91; and Vasquez et al., "METH-1, a human ortholog of ADAMTS-1, and METH-2 are members of a new family of proteins with angio-inhibitory activity," J. Biol. Chem. (Aug. 1999) 274:23349-57.

20

The present invention is directed to novel metalloproteases having thrombospondin domain(s) (MPTS proteins) and polypeptides related thereto, as well as nucleic acid compositions encoding the same, are provided. The subject polypeptide and nucleic acid compositions find use in a variety of applications, including diagnostic

25 applications, therapeutic agent screening applications, as well as therapeutic applications for a variety of different conditions. Also provided are methods of treating disease conditions associated with aggrecanase activity, e.g. conditions characterized by the presence of aggrecan cleavage products, such as rheumatoid- and osteo-arthritis.

MPTS protein of the subject invention. Figure 1B provides the amino acid sequence

of MPTS-15. Figure 1C provides an alignment of the amino acid sequence of the subject MPTS-15 with the amino acid sequence of ADAMTS-6, a sequence disclosed in Hurskainen et al., J. Biol. Chem. (Sept. 1999) 274: 25555-25563.

Figure 2A provides the sequence of a nucleic acid that encodes MPTS-10, an MPTS protein of the subject invention. Figure 2B provides the amino acid sequence of MPTS-10.

Figure 3A provides the sequence of a nucleic acid that encodes MPTS-19, an MPTS protein of the subject invention. Figure 3B provides the amino acid sequence of MPTS-19.

Figure 4A provides the sequence of a nucleic acid that encodes MPTS-20, an MPTS protein of the subject invention. Figure 4B provides the amino acid sequence of MPTS-20.

Novel MPTS proteins and polypeptides related thereto, as well as nucleic acid compositions encoding the same, are provided. The subject polypeptide and/or nucleic acid compositions find use in a variety of different applications, including research, diagnostic, and therapeutic agent screening/discovery/ preparation applications. Also provided are methods of treating disease conditions associated with MPTS, including aggrecanase, function, e.g. diseases characterized by the presence of aggrecan cleavage products, such as rheumatoid- and osteo-arthritis.

Novel metalloproteases having thrombospondin domain(s) (also known as MPTS proteins, ADAMTS proteins or aggrecanase proteins), as well as polypeptide compositions related thereto, are provided. The term polypeptide composition as used herein refers to both the full length protein, as well as portions or fragments thereof. Also included in this term are variations of the naturally occurring human protein, where such variations are homologous or substantially similar to the naturally occurring protein, as described in greater detail below. In the following description of the subject invention, the term "MPTS" is used to refer not only to the specific human MPTS proteins disclosed herein (i.e. MPTS-10; MPTS-15; MPTS-19 and MPTS-20), but also to homologs thereof expressed in non-human species, e.g. murine, rat and other mammalian species.



Specific human MPTS proteins of interest are MPTS-15, MPTS-10, MPTS-19 and MPTS-20. MPTS-15 has an amino acid sequence as shown in Fig. 1B and identified as SEQ ID NO:01. MPTS-10 has an amino acid sequence as shown in Fig. 2B and identified as SEQ ID NO:03. MPTS-19 has an amino acid sequence as shown in Fig. 3B and identified as SEQ ID NO:05. MPTS-20 has an amino acid sequence as shown in Fig. 4B and identified as SEQ ID NO:07. The subject MPTS proteins have a molecular weight based on their amino acid sequence of at least about 90 kDal, where the molecular weight based on the amino acid sequence may be substantially higher in certain embodiments. The true molecular weight of the subject MPTS proteins may vary due to glycosylation and/or other posttranslational modifications.

Also provided by the subject invention are MPTS polypeptide compositions. The term polypeptide composition as used herein refers to both the full length proteins as well as portions or fragments thereof. Also included in this term are variations of the naturally occurring proteins, where such variations are homologous or substantially similar to the naturally occurring protein, be the naturally occurring protein the human protein, mouse protein, or protein from some other species which naturally expresses an MPTS protein, usually a mammalian species. A candidate homologous protein is substantially similar to an MPTS protein of the subject invention, and therefore is an MPTS protein of the subject invention, if the candidate protein has a sequence that has at least about 35%, usually at least about 45% and more usually at least about 60% sequence identity with an MPTS protein, as determined using MegAlign, DNASTAR (1998) clustal algorithm as described in D. G. Higgins and P.M. Sharp, "Fast and Sensitive multiple Sequence Alignments on a Microcomputer," (1989) CABIOS, 5: 151-153. (Parameters used are ktuple 1, gpa penalty 3, window, 5 and diagonals saved 5). In the following description of the subject invention, the term "MPTS-protein" is used to refer not only to the human MPTS proteins, but also to homologs thereof expressed in non-human species, e.g. murine, rat and other mammalian species.

usually at least about 65%, and more usually at least about 70%. In many preferred

embodiments, the sequence identity is at least about 90%, usually at least about 95% and more usually at least about 99% over the entire length of the protein.

In many embodiments, the proteins of the subject invention are enzymes,  
5 particularly proteinases and more particularly a metalloproteinases. The subject proteins of this embodiment are characterized by having aggrecanase activity. As such, the subject proteins are capable of cleaving aggrecan in an interglobular domain, particularly between the G1 and G2 domains, and more particularly at the Glu<sup>373</sup>-Ala<sup>374</sup> bond of human aggrecan, to produce a cleavage product having an N-terminal sequence  
10 of ARGSVIL.

In addition to the proteins described above, homologs or proteins (or fragments thereof) from other species, i.e. other animal or plant species, are also provided, where such homologs or proteins may be from a variety of different types of species, usually  
15 mammals, e.g. rodents, such as mice, rats; domestic animals, e.g. horse, cow, dog, cat; and humans. By homolog is meant a protein having at least about 35 %, usually at least about 40% and more usually at least about 60 % amino acid sequence identity with one of the specific human MPTS proteins as identified above (i.e. with a protein having the amino acid sequence of SEQ ID NOS:01, 03, 05 or 07), where sequence identity is  
20 determined as described *supra*.

The proteins of the subject invention are present in a non-naturally occurring environment, e.g. they are separated from their naturally occurring environment. In certain embodiments, the subject proteins are present in a composition that is enriched  
25 for the subject protein as compared to its naturally occurring environment. For example, purified protein is provided, where by purified is meant that the protein is present in a composition that is substantially free of non-MPTS proteins, where by substantially free is meant that less than 90 %, usually less than 60 % and more usually less than 50 % of the composition is made up of non- MPTS proteins. The proteins of  
30 the subject invention may also be present as an isolate, by which is meant that the protein is substantially free of other proteins and other naturally occurring biologic molecules, such as oligosaccharides, polynucleotides and fragments thereof, and the like, where the term "substantially free" in this instance means that less than 70 %,

usually less than 60% and more usually less than 50 % of the composition containing the isolated protein is some other naturally occurring biological molecule. In certain embodiments, the proteins are present in substantially pure form, where by "substantially pure form" is meant at least 95%, usually at least 97% and more usually at least 99% pure.

In addition to the naturally occurring proteins, polypeptides which vary from the naturally occurring proteins are also provided, e.g. MPTS polypeptides. By MPTS polypeptide is meant an amino acid sequence encoded by an open reading frame (ORF) of the gene encoding the MPTS, described in greater detail below, including the full length protein and fragments thereof, particularly biologically active fragments and/or fragments corresponding to functional domains, e.g. protease domain, thrombospondin domain, and the like; and including fusions of the subject polypeptides to other proteins or parts thereof. Fragments of interest will typically be at least about 10 aa in length, usually at least about 50 aa in length, and may be as long as 300 aa in length or longer, but will usually not exceed about 1000 aa in length, where the fragment will have a stretch of amino acids that is identical to the subject protein of at least about 10 aa, and usually at least about 15 aa, and in many embodiments at least about 50 aa in length. Where the fragment is an MPTS-15 fragment, it preferably includes at least a substantial portion of the protease domain of the wild type protein, where by substantial amount is at least 50%, usually at least 60 % and more usually at least 70 % of the sequence of this domain of the MPTS-15 protein. For example, the MPTS-15 fragment generally includes a sequence which upon alignment with the sequence of residues from the protease domain of the wild type sequence, shows an identity with the aligned region of the wild type sequence of this domain of at least about 50%, usually at least about 60% and more usually at least about 70%, wherein in many embodiments the percent identity may be much higher, e.g. 75, 80, 85, 90 or 95% or higher, e.g. 99%.

The subject proteins may be derived from:

1. naturally occurring sources, e.g. from animal sources, e.g. cartilage and the like. The subject proteins may also be derived from synthetic means,

e.g. by expressing a recombinant gene encoding protein of interest in a suitable host, as described in greater detail below. Any convenient protein purification procedures may be employed, where suitable protein purification methodologies are described in Guide to Protein Purification, (Deuthser ed.) (Academic Press, 1990). For example, a lysate  
5 may prepared from the original source, e.g. chondrocytes or the expression host, and purified using HPLC, exclusion chromatography, gel electrophoresis, affinity chromatography, and the like.

Also provided are nucleic acid compositions encoding MPTS proteins or  
10 fragments thereof, as well as the MPTS homologues of the present invention. By nucleic acid composition is meant a composition comprising a sequence of DNA having an open reading frame that encodes an MPTS polypeptide of the subject invention, i.e. an *mpts* gene, and is capable, under appropriate conditions, of being expressed as MPTS. Also encompassed in this term are nucleic acids that are homologous or substantially  
15 similar or identical to the nucleic acids encoding MPTS proteins. Thus, the subject invention provides genes encoding the human MPTS proteins of the subject invention and homologs thereof. The human MPTS15 gene is shown in Fig. 1A, where the sequence shown in Fig. 1A is identified as SEQ ID NO:02, *infra*. The human MPTS10 gene is shown in Fig. 2A, where the sequence shown in Fig. 2A is identified as SEQ ID  
20 NO:04, *infra*. The human MPTS19 gene is shown in Fig. 3A, where the sequence shown in Fig. 3A is identified as SEQ ID NO:06, *infra*. The human MPTS20 gene is shown in Fig. 4A, where the sequence shown in Fig. 4A is identified as SEQ ID NO:08, *infra*.

The source of homologous genes may be any species, e.g., primate species,  
25 particularly human; rodents, such as rats and mice, canines, felines, bovines, ovines, equines, yeast, nematodes, etc. Between mammalian species, e.g., human and mouse, homologs have substantial sequence similarity, e.g. at least 75% sequence identity, usually at least 90%, more usually at least 95% between nucleotide sequences. Sequence similarity is calculated based on a reference sequence, which may be a subset of a larger  
30 sequence, such as a conserved motif, coding region, flanking region, etc. A reference sequence will usually be at least about 18 nt long, more usually at least about 30 nt long, and may extend to the complete sequence that is being compared. Algorithms for sequence analysis are known in the art, such as BLAST, described in Altschul *et al.*

(1990), *J. Mol. Biol.* 215:403-10 (using default settings, i.e. parameters  $w=4$  and  $T=17$ ).

The sequences provided herein are essential for recognizing MPTS-, including aggrecanase-, related and homologous proteins, and the nucleic acids encoding the same, in database searches. Of particular interest in certain embodiments are nucleic acids of substantially the same length as the nucleic acids identified as SEQ ID NO:02, 04, 06 and 08 and have sequence identity to one of these sequences of at least about 90%, usually at least about 95% and more usually at least about 99% over the entire length of the nucleic acid.

10 Nucleic acids encoding the proteins and polypeptides of the subject invention may be cDNA or genomic DNA or a fragment thereof. The term "MPTS gene" shall be intended to mean the open reading frame encoding specific MPTS proteins and polypeptides, and introns, as well as adjacent 5' and 3' non-coding nucleotide sequences involved in the regulation of expression, up to about 20 kb beyond the coding  
15 region, but possibly further in either direction. The gene may be introduced into an appropriate vector for extrachromosomal maintenance or for integration into a host genome.

The term "cDNA" as used herein is intended to include all nucleic acids that  
20 share the arrangement of sequence elements found in native mature mRNA species, where sequence elements are exons and 5' and 3' non-coding regions. Normally mRNA species have contiguous exons, with the intervening introns, when present, being removed by nuclear RNA splicing, to create a continuous open reading frame encoding an MPTS protein.

25 A genomic sequence of interest comprises the nucleic acid present between the initiation codon and the stop codon, as defined in the listed sequences, including all of the introns that are normally present in a native chromosome. It may further include 5' and 3' untranslated regions found in the original DNA. It may further include, in addition

either the 5' or 3' end of the transcribed region. The genomic DNA may be isolated as a

fragment of 100 kbp or smaller; and substantially free of flanking chromosomal sequence. The genomic DNA flanking the coding region, either 3' or 5', or internal regulatory sequences as sometimes found in introns, contains sequences required for proper tissue and stage specific expression.

5

The nucleic acid compositions of the subject invention may encode all or a part of the subject MPTS protein. Double or single stranded fragments may be obtained from the DNA sequence by chemically synthesizing oligonucleotides in accordance with conventional methods, by restriction enzyme digestion, by PCR amplification, *etc.* For  
10 the most part, DNA fragments will be of at least 15 nt, usually at least 18 nt or 25 nt, and may be at least about 50 nt.

The subject genes are isolated and obtained in substantial purity, generally as other than an intact chromosome. Usually, the DNA will be obtained substantially free  
15 of other nucleic acid sequences that do not include an MPTS gene sequence or fragment thereof, generally being at least about 50%, usually at least about 90% pure and are typically "recombinant", *i.e.* flanked by one or more nucleotides with which it is not normally associated on a naturally occurring chromosome.

20 In addition to the plurality of uses described in greater detail in following sections, the subject nucleic acid compositions find use in the preparation of all or a portion of the MPTS polypeptides, as described above. The provided polynucleotide (*e.g.*, a polynucleotide having a sequence of SEQ ID NO:02, 04, 06 or 08), the corresponding cDNA, or the full-length gene is used to express a partial or complete  
25 gene product. Constructs of polynucleotides having a sequences of SEQ ID NOs: 02, 04, 06 or 08 can be generated synthetically. Alternatively, single-step assembly of a gene and entire plasmid from large numbers of oligodeoxyribonucleotides is described by, *e.g.*, Stemmer *et al.*, *Gene (Amsterdam)* (1995) 164(1):49-53. In this method, assembly PCR (the synthesis of long DNA sequences from large numbers of oligodeoxy-  
30 ribonucleotides (oligos)) is described. The method is derived from DNA shuffling (Stemmer, *Nature* (1994) 370:389-391), and does not rely on DNA ligase, but instead relies on DNA polymerase to build increasingly longer DNA fragments during the assembly process. Appropriate polynucleotide constructs are purified using standard

recombinant DNA techniques as described in, for example, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Ed., (1989) Cold Spring Harbor Press, Cold Spring Harbor, NY, and under current regulations described in United States Dept. of HHS, National Institute of Health (NIH) Guidelines for Recombinant DNA Research.

5

Polynucleotide molecules comprising a polynucleotide sequence provided herein are propagated by placing the molecule in a vector. Viral and non-viral vectors are used, including plasmids. The choice of plasmid will depend on the type of cell in which propagation is desired and the purpose of propagation. Certain vectors are useful for  
10 amplifying and making large amounts of the desired DNA sequence. Other vectors are suitable for expression in cells in culture. Still other vectors are suitable for transfer and expression in cells in a whole animal or person. The choice of appropriate vector is well within the skill of the art. Many such vectors are available commercially. The partial or full-length polynucleotide is inserted into a vector typically by means of DNA ligase  
15 attachment to a cleaved restriction enzyme site in the vector. Alternatively, the desired nucleotide sequence can be inserted by homologous recombination *in vivo*. Typically this is accomplished by attaching regions of homology to the vector on the flanks of the desired nucleotide sequence. Regions of homology are added by ligation of oligonucleotides, or by polymerase chain reaction using primers comprising both the  
20 region of homology and a portion of the desired nucleotide sequence, for example.

For expression, an expression cassette or system may be employed. The gene product encoded by a polynucleotide of the invention is expressed in any convenient expression system, including, for example, bacterial, yeast, insect, amphibian and  
25 mammalian systems. Suitable vectors and host cells are described in U.S. Patent No. 5,654,173. In the expression vector, an MPIS encoding polynucleotide, e.g. as set forth in SEQ ID NO: 02, 04, 06 or 08, is linked to a regulatory sequence as appropriate to obtain the desired expression properties. These can include promoters (attached either at the 5' end of the sense strand or at the 3' end of the antisense strand), enhancers,  
30 terminators, and other regulatory elements. The regulatory sequence can be directly  
35 attached to the desired nucleotide sequence using the techniques described above for

linkage to vectors. Any techniques known in the art can be used. In other words, the expression vector will provide a transcriptional and translational initiation region, which may be inducible or constitutive, where the coding region is operably linked under the transcriptional control of the transcriptional initiation region, and a  
5 transcriptional and translational termination region. These control regions may be native to the subject MPTS gene, or may be derived from exogenous sources.

Expression vectors generally have convenient restriction sites located near the promoter sequence to provide for the insertion of nucleic acid sequences encoding  
10 heterologous proteins. A selectable marker operative in the expression host may be present. Expression vectors may be used for the production of fusion proteins, where the exogenous fusion peptide provides additional functionality, i.e. increased protein synthesis, stability, reactivity with defined antisera, an enzyme marker, e.g.  $\beta$ -galactosidase, etc.

15 Expression cassettes may be prepared comprising a transcription initiation region, the gene or fragment thereof, and a transcriptional termination region. Of particular interest is the use of sequences that allow for the expression of functional epitopes or domains, usually at least about 8 amino acids in length, more usually at least  
20 about 15 amino acids in length, to about 25 amino acids, and up to the complete open reading frame of the gene. After introduction of the DNA, the cells containing the construct may be selected by means of a selectable marker, the cells expanded and then used for expression.

25 The MPTS proteins and polypeptides may be expressed in prokaryotes or eukaryotes in accordance with conventional ways, depending upon the purpose for expression. For large scale production of the protein, a unicellular organism, such as *E. coli*, *B. subtilis*, *S. cerevisiae*, insect cells in combination with baculovirus vectors, or cells of a higher organism such as vertebrates, particularly mammals, e.g. COS 7 cells, HEK  
30 293, CHO, *Xenopus* Oocytes, etc., may be used as the expression host cells. In some situations, it is desirable to express the gene in eukaryotic cells, where the expressed protein will benefit from native folding and post-translational modifications. Small peptides can also be synthesized in the laboratory. Polypeptides that are subsets of the



complete protein sequence may be used to identify and investigate parts of the protein important for function.

Specific expression systems of interest include bacterial, yeast, insect cell and  
 5 mammalian cell derived expression systems. Representative systems from each of these categories is are provided below

Bacteria. Expression systems in bacteria include those described in Chang *et al.*,  
*Nature* (1978) 275:615; Goeddel *et al.*, *Nature* (1979) 281:544; Goeddel *et al.*, *Nucleic*  
 10 *Acids Res.* (1980) 8:4057; EP 0 036,776; U.S. Patent No. 4,551,433; DeBoer *et al.*, *Proc.*  
*Natl. Acad. Sci. (USA)* (1983) 80:21-25; and Siebenlist *et al.*, *Cell* (1980) 20:269.

Yeast. Expression systems in yeast include those described in Hinnen *et al.*, *Proc.*  
*Natl. Acad. Sci. (USA)* (1978) 75 1929; Ito *et al.*, *J. Bacteriol.* (1983) 153:163; Kurtz *et al.*,  
*Mol. Cell. Biol.* (1986) 6 142; Kunze *et al.*, *J. Basic Microbiol.* (1985) 25:141; Gleeson *et*  
 15 *al.*, *J. Gen. Microbiol.* (1986) 132 3459; Roggenkamp *et al.*, *Mol. Gen. Genet.* (1986)  
 202:302; Das *et al.*, *J. Bacteriol.* (1984) 158:1165; De Louvencourt *et al.*, *J. Bacteriol.*  
 (1983) 154:737; Van den Berg *et al.*, *Bio/Technology* (1990) 8:135; Kunze *et al.*, *J. Basic*  
*Microbiol.* (1985) 25:141; Cregg *et al.*, *Mol. Cell. Biol.* (1985) 5:3376; U.S. Patent Nos.  
 4,837,148 and 4,929,555; Beach and Nurse, *Nature* (1981) 300:706; Davidow *et al.*, *Curr.*  
 20 *Genet.* (1985) 10:380; Gaillardin *et al.*, *Curr. Genet.* (1985) 10:49; Ballance *et al.*,  
*Biochem. Biophys. Res. Commun.* (1983) 112:284-289; Tilburn *et al.*, *Gene* (1983)  
 26:205-221; Yelton *et al.*, *Proc. Natl. Acad. Sci. (USA)* (1984) 81:1470-1474; Kelly and  
 Hynes, *EMBO J.* (1985) 4:475-479; EP 0 244,234; and WO 91/00357.

Insect Cells. Expression of heterologous genes in insects is accomplished as  
 25 described in U.S. Patent No. 4,745,051; Friesen *et al.*, "The Regulation of Baculovirus  
 Gene Expression", in: *The Molecular Biology Of Baculoviruses* (1986) (W. Doertler, ed.);  
 EP 0 127,839; EP 0 155,476; and Vlak *et al.*, *J. Gen. Virol.* (1988) 69:765-776; Miller *et al.*,  
*Ann. Rev. Microbiol.* (1988) 42:177; Carbonell *et al.*, *Gene* (1988) 73:409; Maeda *et al.*,  
*Nature* (1985) 315:592-594; Lebacqz-Verheyden *et al.*, *Mol. Cell. Biol.* (1988) 8:3129;  
 30 Suckiyaporn, *Dev. Mol. Biol. Sci. (USA)* (1987) 8:884-890; and Vlak *et al.*, *J. Gen. Virol.* (1988)

*al.*, *Bio/Technology* (1988) 6:47-55, Miller *et al.*, *Generic Engineering* (1986) 8:277-279, and Maeda *et al.*, *Nature* (1985) 315:592-594.

Mammalian Cells. Mammalian expression is accomplished as described in Dijkema *et al.*, *EMBO J.* (1985) 4:761, Gorman *et al.*, *Proc. Natl. Acad. Sci. (USA)* (1982) 79:6777, Boshart *et al.*, *Cell* (1985) 41:521 and U.S. Patent No. 4,399,216. Other features of mammalian expression are facilitated as described in Ham and Wallace, *Meth. Enz.* (1979) 58:44, Barnes and Sato, *Anal. Biochem.* (1980) 102:255, U.S. Patent Nos. 4,767,704, 4,657,866, 4,927,762, 4,560,655, WO 90/103430, WO 87/00195, and U.S. RE 30,985.

10

When any of the above host cells, or other appropriate host cells or organisms, are used to replicate and/or express the polynucleotides or nucleic acids of the invention, the resulting replicated nucleic acid, RNA, expressed protein or polypeptide, is within the scope of the invention as a product of the host cell or organism. The product is recovered by any appropriate means known in the art.

15

Once the gene corresponding to a selected polynucleotide is identified, its expression can be regulated in the cell to which the gene is native. For example, an endogenous gene of a cell can be regulated by an exogenous regulatory sequence as disclosed in U.S. Patent No. 5,641,670.

20

The subject polypeptide and nucleic acid compositions find use in a variety of different applications, including general applications, diagnostic applications, and therapeutic agent screening/discovery; preparation applications, as well as in therapeutic compositions and methods employing the same.

25

The subject nucleic acid compositions find use in a variety of general applications. General applications of interest include: the identification of MPTS homologs; as a source of novel promoter elements; the identification of MPTS expression regulatory factors; as probes and primers in hybridization applications, e.g. PCR; the identification of expression patterns in biological specimens; the preparation of cell or animal models for MPTS function; the preparation of *in vitro* models for MPTS function; etc.

30

- Homologs of the subject genes are identified by any of a number of methods. A fragment of the provided cDNA may be used as a hybridization probe against a cDNA library from the target organism of interest, where low stringency conditions are used.
- 5 The probe may be a large fragment, or one or more short degenerate primers. Nucleic acids having sequence similarity are detected by hybridization under low stringency conditions, for example, at 50°C and 6xSSC (0.9 M sodium chloride/0.09 M sodium citrate) and remain bound when subjected to washing at 55°C in 1xSSC (0.15 M sodium chloride/0.015 M sodium citrate). Sequence identity may be determined by
- 10 hybridization under stringent conditions, for example, at 50°C or higher and 0.1xSSC (15 mM sodium chloride/0.15 mM sodium citrate). Nucleic acids having a region of substantial identity to the provided sequences, e.g. allelic variants, genetically altered versions of the gene, etc., bind to the provided sequences under stringent hybridization conditions. By using probes, particularly labeled probes of DNA sequences, one can
- 15 isolate homologous or related genes.

- The sequence of the 5' flanking region may be utilized for promoter elements, including enhancer binding sites, that provide for developmental regulation in tissues where the subject MPFS gene is expressed. The tissue specific expression is useful for
- 20 determining the pattern of expression, and for providing promoters that mimic the native pattern of expression. Naturally occurring polymorphisms in the promoter region are useful for determining natural variations in expression, particularly those that may be associated with disease.

- 25 Alternatively, mutations may be introduced into the promoter region to determine the effect of altering expression in experimentally defined systems. Methods for the identification of specific DNA motifs involved in the binding of transcriptional factors are known in the art, e.g. sequence similarity to known binding motifs, gel retardation studies, etc. For examples, see Blackwell *et al.* (1995), *Mol. Med.* 1:194-205;

The regulatory sequences may be used to identify *cis* acting sequences required for transcriptional or translational regulation of MPTS gene expression, especially in different tissues or stages of development, and to identify *cis* acting sequences and *trans*-acting factors that regulate or mediate MPTS gene expression. Such transcription or  
5 translational control regions may be operably linked to an MPTS gene in order to promote expression of wild type or altered MPTS or other proteins of interest in cultured cells, or in embryonic, fetal or adult tissues, and for gene therapy.

Small DNA fragments are useful as primers for PCR, hybridization screening  
10 probes, *etc.* Larger DNA fragments, *i.e.* greater than 100 nt are useful for production of the encoded polypeptide, as described in the previous section. For use in geometric amplification reactions, such as geometric PCR, a pair of primers will be used. The exact composition of the primer sequences is not critical to the invention, but for most applications the primers will hybridize to the subject sequence under stringent  
15 conditions, as known in the art. It is preferable to choose a pair of primers that will generate an amplification product of at least about 50 nt, preferably at least about 100 nt. Algorithms for the selection of primer sequences are generally known, and are available in commercial software packages. Amplification primers hybridize to complementary strands of DNA, and will prime towards each other.

20

The DNA may also be used to identify expression of the gene in a biological specimen. The manner in which one probes cells for the presence of particular nucleotide sequences, as genomic DNA or RNA, is well established in the literature. Briefly, DNA or mRNA is isolated from a cell sample. The mRNA may be amplified by  
25 RT-PCR, using reverse transcriptase to form a complementary DNA strand, followed by polymerase chain reaction amplification using primers specific for the subject DNA sequences. Alternatively, the mRNA sample is separated by gel electrophoresis, transferred to a suitable support, *e.g.* nitrocellulose, nylon, *etc.*, and then probed with a fragment of the subject DNA as a probe. Other techniques, such as oligonucleotide  
30 ligation assays, *in situ* hybridizations, and hybridization to DNA probes arrayed on a solid chip may also find use. Detection of mRNA hybridizing to the subject sequence is indicative of MPTS gene expression in the sample.

The sequence of an MPTS gene, including flanking promoter regions and coding regions, may be mutated in various ways known in the art to generate targeted changes in promoter strength, sequence of the encoded protein, *etc.* The DNA sequence or protein product of such a mutation will usually be substantially similar to the sequences provided herein, *i.e.* will differ by at least one nucleotide or amino acid, respectively, and may differ by at least two but not more than about ten nucleotides or amino acids. The sequence changes may be substitutions, insertions, deletions, or a combination thereof. Deletions may further include larger changes, such as deletions of a domain or exon. Other modifications of interest include epitope tagging, *e.g.* with the FLAG system, HA, *etc.* For studies of subcellular localization, fusion proteins with green fluorescent proteins (GFP) may be used.

Techniques for *in vitro* mutagenesis of cloned genes are known. Examples of protocols for site specific mutagenesis may be found in Gustin *et al.* (1993), *Biotechniques* 14:22; Barany (1985), *Gene* 37:111-23. Colicelli *et al.* (1985), *Mol. Gen. Genet.* 199:537-9; and Prentki *et al.* (1984), *Gene* 29:303-13. Methods for site specific mutagenesis can be found in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, CSH Press 1989, pp. 15.3-15.108; Weiner *et al.* (1993), *Gene* 126:35-41; Sayers *et al.* (1992), *Biotechniques* 13:592-6; Jones and Winistorfer (1992), *Biotechniques* 12:528-30; Barton *et al.* (1990), *Nucleic Acids Res* 18:7349-55; Marotti and Tomich (1989), *Gene Anal. Tech.* 6:67-70; and Zhu (1989), *Anal Biochem* 177:120-4. Such mutated genes may be used to study structure-function relationships of an MPTS protein, or to alter properties of the protein that affect its function or regulation.

The subject nucleic acids can be used to generate transgenic, non human animals or site specific gene modifications in cell lines. Transgenic animals may be made through homologous recombination, where the endogenous locus is altered. Alternatively, a nucleic acid construct is randomly integrated into the genome. Vectors for stable integration include plasmids, retroviruses and other animal viruses, YACs,

regulation. Of interest is the use of the subject genes to construct transgenic animal

models of MPTS related disease conditions, including aggrecanase related disease conditions, e.g. disease conditions associated with aggrecanase activity, such as arthritis. Thus, transgenic animal models of the subject invention include endogenous MPTS gene knockouts in which expression of endogenous MPTS is at least reduced if not  
5 eliminated, where such animals also typically express an MPTS peptide of the subject invention, e.g. the specific MPTS proteins of the subject invention or a fragment thereof. Where a nucleic acid having a sequence found in the human MPTS gene is introduced, the introduced nucleic acid may be either a complete or partial sequence of the MPTS gene. A detectable marker, such as *lac Z* may be introduced into the MPTS  
10 locus, where upregulation of gene expression will result in an easily detected change in phenotype. One may also provide for expression of the gene or variants thereof in cells or tissues where it is not normally expressed, at levels not normally present in such cells or tissues.

15 DNA constructs for homologous recombination will comprise at least a portion of the an MPTS gene of the subject invention, wherein the gene has the desired genetic modification(s), and includes regions of homology to the target locus. DNA constructs for random integration need not include regions of homology to mediate recombination. Conveniently, markers for positive and negative selection are included.  
20 Methods for generating cells having targeted gene modifications through homologous recombination are known in the art. For various techniques for transfecting mammalian cells, see Keown *et al.* (1990), *Meth. Enzymol.* 185:527-537.

For embryonic stem (ES) cells, an ES cell line may be employed, or embryonic  
25 cells may be obtained freshly from a host, e.g. mouse, rat, guinea pig, *etc.* Such cells are grown on an appropriate fibroblast-feeder layer or grown in the presence of leukemia inhibiting factor (LIF). When ES or embryonic cells have been transformed, they may be used to produce transgenic animals. After transformation, the cells are plated onto a feeder layer in an appropriate medium. Cells containing the construct may be detected  
30 by employing a selective medium. After sufficient time for colonies to grow, they are picked and analyzed for the occurrence of homologous recombination or integration of the construct. Those colonies that are positive may then be used for embryo manipulation and blastocyst injection. Blastocysts are obtained from 4 to 6 week old

superovulated females. The ES cells are trypsinized, and the modified cells are injected into the blastocoel of the blastocyst. After injection, the blastocysts are returned to each uterine horn of pseudopregnant females. Females are then allowed to go to term and the resulting offspring screened for the construct. By providing for a different  
5 phenotype of the blastocyst and the genetically modified cells, chimeric progeny can be readily detected.

The chimeric animals are screened for the presence of the modified gene and males and females having the modification are mated to produce homozygous progeny.  
10 If the gene alterations cause lethality at some point in development, tissues or organs can be maintained as allogeneic or congenic grafts or transplants, or in *in vitro* culture. The transgenic animals may be any non-human mammal, such as laboratory animals, domestic animals, etc. The transgenic animals may be used in functional studies, drug screening, etc., e.g. to determine the effect of a candidate drug on aggrecanase activity.

15 Also provided are methods of diagnosing disease states based on observed levels of an MPTS protein or the expression level of the gene in a biological sample of interest. Samples, as used herein, include biological fluids such as blood, cerebrospinal fluid, tears, saliva, lymph, dialysis fluid, and the like; organ or tissue culture derived fluids;  
20 and fluids extracted from physiological tissues. Also included in the term are derivatives and fractions of such fluids. The cells may be dissociated, in the case of solid tissues, or tissue sections may be analyzed. Alternatively a lysate of the cells may be prepared.

A number of methods are available for determining the expression level of a  
25 gene or protein in a particular sample. Diagnosis may be performed by a number of methods to determine the absence or presence or altered amounts of normal or abnormal MPTS in a patient sample. For example, detection may utilize staining of cells or histological sections with labeled antibodies, performed in accordance with conventional methods. Cells are permeabilized to stain cytoplasmic molecules. The

other labels for direct detection. Alternatively, a second stage antibody or reagent is

used to amplify the signal. Such reagents are well known in the art. For example, the primary antibody may be conjugated to biotin, with horseradish peroxidase-conjugated avidin added as a second stage reagent. Alternatively, the secondary antibody conjugated to a fluorescent compound, e.g. fluorescein, rhodamine, Texas red, *etc.* Final  
5 detection uses a substrate that undergoes a color change in the presence of the peroxidase. The absence or presence of antibody binding may be determined by various methods, including flow cytometry of dissociated cells, microscopy, radiography, scintillation counting, *etc.*

10 Alternatively, one may focus on the expression of the MPTS gene. Biochemical studies may be performed to determine whether a sequence polymorphism in an MPTS coding region or control regions is associated with disease. Disease associated polymorphisms may include deletion or truncation of the gene, mutations that alter expression level, that affect the activity of the protein, *etc.*

15 Changes in the promoter or enhancer sequence that may affect expression levels of MPTS can be compared to expression levels of the normal allele by various methods known in the art. Methods for determining promoter or enhancer strength include quantitation of the expressed natural protein; insertion of the variant control element  
20 into a vector with a reporter gene such as  $\beta$ -galactosidase, luciferase, chloramphenicol acetyltransferase, *etc.* that provides for convenient quantitation; and the like.

A number of methods are available for analyzing nucleic acids for the presence of a specific sequence, e.g. a disease associated polymorphism. Where large amounts of  
25 DNA are available, genomic DNA is used directly. Alternatively, the region of interest is cloned into a suitable vector and grown in sufficient quantity for analysis. Cells that express an MPTS protein may be used as a source of mRNA, which may be assayed directly or reverse transcribed into cDNA for analysis. The nucleic acid may be amplified by conventional techniques, such as the polymerase chain reaction (PCR), to  
30 provide sufficient amounts for analysis. The use of the polymerase chain reaction is described in Saiki, *et al.* (1985), *Science* 239:487, and a review of techniques may be found in Sambrook, *et al.* Molecular Cloning: A Laboratory Manual, CSH Press 1989, pp.14.2-14.33. Alternatively, various methods are known in the art that utilize



oligonucleotide ligation as a means of detecting polymorphisms, for examples see Riley *et al.* (1990), *Nucl. Acids Res.* 18:2887-2890; and Delahunty *et al.* (1996), *Am. J. Hum. Genet.* 58:1239-1246.

5           A detectable label may be included in an amplification reaction. Suitable labels include fluorochromes, *e.g.* fluorescein isothiocyanate (FITC), rhodamine, Texas Red, phycoerythrin, allophycocyanin, 6-carboxyfluorescein (6-FAM), 2',7'-dimethoxy-4',5'-dichloro-6-carboxyfluorescein (JOE), 6-carboxy-X-rhodamine (ROX), 6-carboxy-2',4',7',4,7-hexachlorofluorescein (HEX), 5-carboxyfluorescein (5-FAM) or N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA), radioactive labels, *e.g.*  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^3\text{H}$ ; *etc.* The label may be a two stage system, where the amplified DNA is conjugated to biotin, haptens, *etc.* having a high affinity binding partner, *e.g.* avidin, specific antibodies, *etc.*, where the binding partner is conjugated to a detectable label. The label may be conjugated to one or both of the primers. Alternatively, the pool of nucleotides used in  
10           the amplification is labeled, so as to incorporate the label into the amplification  
15           product.

          The sample nucleic acid, *e.g.* amplified or cloned fragment, is analyzed by one of a number of methods known in the art. The nucleic acid may be sequenced by dideoxy  
20           or other methods, and the sequence of bases compared to a wild-type gene sequence. Hybridization with the variant sequence may also be used to determine its presence, by Southern blots, dot blots, *etc.* The hybridization pattern of a control and variant sequence to an array of oligonucleotide probes immobilized on a solid support, as described in US 5,445,934, or in WO 95/35505, may also be used as a means of detecting  
25           the presence of variant sequences. Single strand conformational polymorphism (SSCP) analysis, denaturing gradient gel electrophoresis (DGGE), and heteroduplex analysis in gel matrices are used to detect conformational changes created by DNA sequence variation as alterations in electrophoretic mobility. Alternatively, where a polymorphism creates or destroys a recognition site for a restriction endonuclease, the

Screening for mutations in MPTS may be based on the functional or antigenic characteristics of the protein. Protein truncation assays are useful in detecting deletions that may affect the biological activity of the protein. Various immunoassays designed to detect polymorphisms in MPTS proteins may be used in screening. Where many  
5 diverse genetic mutations lead to a particular disease phenotype, functional protein assays have proven to be effective screening tools. The activity of the encoded MPTS protein may be determined by comparison with the wild-type protein.

Diagnostic methods of the subject invention in which the level of MPTS gene  
10 expression is of interest will typically involve comparison of the MPTS nucleic acid abundance of a sample of interest with that of a control value to determine any relative differences, where the difference may be measured qualitatively and/or quantitatively, which differences are then related to the presence or absence of an abnormal MPTS gene expression pattern. A variety of different methods for determine the nucleic acid  
15 abundance in a sample are known to those of skill in the art, where particular methods of interest include those described in: Pietu et al., *Genome Res.* (June 1996) 6: 492-503; Zhao et al., *Gene* (April 24, 1995) 156: 207-213; Soares, *Curr. Opin. Biotechnol.* (October 1997) 8: 542-546; Raval, *J. Pharmacol Toxicol Methods* (November 1994) 32: 125-127; Chalifour et al., *Anal. Biochem* (February 1, 1994) 216: 299-304; Stolz & Tuan,  
20 *Mol. Biotechnol.* (December 1996) 6: 225-230; Hong et al., *Bioscience Reports* (1982) 2: 907; and McGraw, *Anal. Biochem.* (1984) 143: 298. Also of interest are the methods disclosed in WO 97/27317, the disclosure of which is herein incorporated by reference.

The subject polypeptides find use in various screening assays designed to  
25 identify therapeutic agents. In vitro screening assays can be employed in which the activity of an MPTS polypeptide, e.g. the aggrecanase activity of an MPTS polypeptide, is assessed in the presence of a candidate therapeutic agent and compared to a control, i.e. the activity in the absence of the candidate therapeutic agent. Activity can be determined in a number of different ways, where activity may generally be determined  
30 as ability to cleave aggrecan or at least a fragment thereof, as well as a recombinant polypeptide, that includes the aggrecanase cleavage site, as described above. Such assays are described in U.S. Patent No. 5,872,209 and WO 99/05921 as well as Arner et al., *J. Biol. Chem.* (March 1999) 274: 6594-6601.

Also of interest in screening assays are non-human transgenic animals that express functional MPTS, where such animals are described above. In many embodiments, the animals lack the corresponding endogenous MPTS. In using such animals for screening applications, a test compound(s) is administered to the animal, and the resultant changes in phenotype, e.g. presence of aggrecan produced by cleavage of the Glu<sup>373</sup>-Ala<sup>374</sup> bond, are compared with a control.

Alternatively, in vitro models of MPTS binding activity may be measured in which binding events between MPTS and candidate MPTS modulatory agents are monitored.

A variety of other reagents may be included in the screening assays, depending on the particular screening protocols employed. These include reagents like salts, neutral proteins, e.g. albumin, detergents, etc that are used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Reagents that improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, *etc.* may be used.

A variety of different candidate therapeutic agents that serve as either MPTS agonists or antagonists may be screened by the above methods. Candidate agents encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 50 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines,

including, but not limited to, nucleic acids and derivatives thereof, or any combination of these synthetic or natural compounds. For example, numerous means are available for

random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides and oligopeptides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically  
5 produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means, and may be used to produce combinatorial libraries. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, *etc.* to produce structural analogs.

10

Of particular interest in many embodiments are screening methods that identify agents that selectively modulate, e.g. inhibit, the subject MPTS enzyme and not other proteases.

15

The nucleic acid compositions of the subject invention also find use as therapeutic agents in situations where one wishes to enhance an MPTS activity in a host. The MPTS genes, gene fragments, or the encoded proteins or protein fragments are useful in gene therapy to treat disorders associated with MPTS defects, including aggrecanase defects. Expression vectors may be used to introduce the gene into a cell.

20

Such vectors generally have convenient restriction sites located near the promoter sequence to provide for the insertion of nucleic acid sequences. Transcription cassettes may be prepared comprising a transcription initiation region, the target gene or fragment thereof, and a transcriptional termination region. The transcription cassettes may be introduced into a variety of vectors, e.g. plasmid; retrovirus, e.g. lentivirus;

25

adenovirus; and the like, where the vectors are able to transiently or stably be maintained in the cells, usually for a period of at least about one day, more usually for a period of at least about several days to several weeks.

30

The gene or protein may be introduced into tissues or host cells by any number of routes, including viral infection, microinjection, or fusion of vesicles. Jet injection may also be used for intramuscular administration, as described by Furth *et al.* (1992), *Anal Biochem* 205:365-368. The DNA may be coated onto gold microparticles, and delivered intradermally by a particle bombardment device, or "gene gun" as described in

the literature (see, for example, Tang *et al.* (1992), *Nature* 356:152-154), where gold microprojectiles are coated with the DNA, then bombarded into skin cells.

5           The subject invention provides methods of modulating MPTS, and in many embodiments aggrecanase, activity in a cell, including methods of increasing MPTS activity (e.g. methods of enhancing ), as well as methods of reducing or inhibiting MPTS activity, e.g. methods of stopping or limiting aggrecan cleavage. In such methods, an effective amount of a modulatory agent is contacted with the cell.

10           Also provided are methods of modulating, including enhancing and inhibiting, MPTS activity in a host. In such methods, an effective amount of active agent that modulates the activity of an MPTS protein *in vivo*, e.g. where the agent usually enhances or inhibits the target MPTS activity, is administered to the host. The active agent may be a variety of different compounds, including a naturally occurring or synthetic small  
15           molecule compound, an antibody, fragment or derivative thereof, an antisense composition, and the like.

            Of particular interest in certain embodiments are agents that reduce MPTS activity, including agents that reduce aggrecanase activity, e.g. aggrecan cleavage, by at  
20           least about 10 fold, usually at least about 20 fold and more usually at least about 25 fold, as measure by the Assay described in Arner *et al.* (1999), *supra*. In many embodiments, of particular interest is the use of compounds that reduce aggrecanase activity by at least 100 fold, as compared to a control.

25           Also of interest is the use of agents that, while providing for reduced MPTS, including aggrecanase, activity, do not substantially reduce the activity of other proteinases, if at all. Thus, the agents in this embodiment are selective inhibitors of MPTS. An agent is considered to be selective if it provides for the above reduced aggrecanase activity, but substantially no reduced activity of at least one other

Naturally occurring or synthetic small molecule compounds of interest include numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 50 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof.

Also of interest as active agents are antibodies that at least reduce, if not inhibit, the target MPTS, e.g. aggrecanase, activity in the host. Suitable antibodies are obtained by immunizing a host animal with peptides comprising all or a portion of the target protein, e.g. MPTS-15, MPTS-19 or MPTS-20. Suitable host animals include mouse, rat, sheep, goat, hamster, rabbit, *etc.* The origin of the protein immunogen may be mouse, human, rat, monkey *etc.* The host animal will generally be a different species than the immunogen, e.g. human MPTS used to immunize mice, *etc.*

The immunogen may comprise the complete protein, or fragments and derivatives thereof. Preferred immunogens comprise all or a part of MPTS, where these residues contain the post-translation modifications, such as glycosylation, found on the native target protein. Immunogens comprising the extracellular domain are produced in a variety of ways known in the art, e.g. expression of cloned genes using conventional recombinant methods, isolation from FHEC, *etc.*

For preparation of polyclonal antibodies, the first step is immunization of the host animal with the target protein, where the target protein will preferably be in substantially pure form, comprising less than about 1% contaminant. The immunogen may comprise the complete target protein, fragments or derivatives thereof. To increase the immune response of the host animal, the target protein may be combined with an adjuvant, where suitable adjuvants include alum, dextran, sulfate, large polymeric

anions, oil & water emulsions, e.g. Freund's adjuvant, Freund's complete adjuvant, and the like. The target protein may also be conjugated to synthetic carrier proteins or synthetic antigens. A variety of hosts may be immunized to produce the polyclonal antibodies. Such hosts include rabbits, guinea pigs, rodents, e.g. mice, rats, sheep, goats, and the like. The target protein is administered to the host, usually intradermally, with an initial dosage followed by one or more, usually at least two, additional booster dosages. Following immunization, the blood from the host will be collected, followed by separation of the serum from the blood cells. The Ig present in the resultant antiserum may be further fractionated using known methods, such as ammonium salt fractionation, DEAE chromatography, and the like.

Monoclonal antibodies are produced by conventional techniques. Generally, the spleen and/or lymph nodes of an immunized host animal provide a source of plasma cells. The plasma cells are immortalized by fusion with myeloma cells to produce hybridoma cells. Culture supernatant from individual hybridomas is screened using standard techniques to identify those producing antibodies with the desired specificity. Suitable animals for production of monoclonal antibodies to the human protein include mouse, rat, hamster, etc. To raise antibodies against the mouse protein, the animal will generally be a hamster, guinea pig, rabbit, etc. The antibody may be purified from the hybridoma cell supernatants or ascites fluid by conventional techniques, e.g. affinity chromatography using MPTS bound to an insoluble support, protein A sepharose, etc. Therefore it is an object of the present invention to provide monoclonal antibodies binding specifically to the MPTS proteins of the present invention, more specifically such antibodies which inhibit aggrecanase activity and such antibodies which are human or humanized ones.

The antibody may be produced as a single chain, instead of the normal multimeric structure. Single chain antibodies are described in Jost *et al.* (1994) *J.B.C.*, 269:26267-73, and others. DNA sequences encoding the variable region of the heavy

retains the specificity and affinity of the original antibody

For *in vivo* use, particularly for injection into humans, it is desirable to decrease the antigenicity of the antibody. An immune response of a recipient against the blocking agent will potentially decrease the period of time that the therapy is effective. Methods of humanizing antibodies are known in the art. The humanized antibody may  
5 be the product of an animal having transgenic human immunoglobulin constant region genes (see for example WO 90/10077 and WO 90/04036). Alternatively, the antibody of interest may be engineered by recombinant DNA techniques to substitute the CH1, CH2, CH3, hinge domains, and/or the framework domain with the corresponding human sequence (see WO 92/02190).

10

The use of Ig cDNA for construction of chimeric immunoglobulin genes is known in the art (Liu *et al.* (1987) P.N.A.S. 84:3439 and (1987) J. Immunol. 139:3521). mRNA is isolated from a hybridoma or other cell producing the antibody and used to produce cDNA. The cDNA of interest may be amplified by the polymerase chain  
15 reaction using specific primers (U.S. Patent Nos. 4,683,195 and 4,683,202). Alternatively, a library is made and screened to isolate the sequence of interest. The DNA sequence encoding the variable region of the antibody is then fused to human constant region sequences. The sequences of human constant regions genes may be found in Kabat *et al.* (1991) Sequences of Proteins of Immunological Interest, N.I.H.  
20 publication no. 91-3242. Human C region genes are readily available from known clones. The choice of isotype will be guided by the desired effector functions, such as complement fixation, or activity in antibody-dependent cellular cytotoxicity. Preferred isotypes are IgG1, IgG3 and IgG4. Either of the human light chain constant regions, kappa or lambda, may be used. The chimeric, humanized antibody is then expressed by  
25 conventional methods.

In yet other embodiments, the antibodies may be fully human antibodies. For example, xenogeneic antibodies which are identical to human antibodies may be employed. By xenogenic human antibodies is meant antibodies that are the same as  
30 human antibodies, i.e. they are fully human antibodies, with exception that they are produced using a non-human host which has been genetically engineered to express human antibodies, e.g. WO 98/50433; WO 98,24893 and WO 99/53049.



Antibody fragments, such as Fv, F(ab')<sub>2</sub> and Fab may be prepared by cleavage of the intact protein, e.g. by protease or chemical cleavage. Alternatively, a truncated gene is designed. For example, a chimeric gene encoding a portion of the F(ab')<sub>2</sub> fragment would include DNA sequences encoding the CH1 domain and hinge region of the H chain, followed by a translational stop codon to yield the truncated molecule.

Consensus sequences of H and L J regions may be used to design oligonucleotides for use as primers to introduce useful restriction sites into the J region for subsequent linkage of V region segments to human C region segments. C region cDNA can be modified by site directed mutagenesis to place a restriction site at the analogous position in the human sequence.

Expression vectors include plasmids, retroviruses, YACs, EBV derived episomes, and the like. A convenient vector is one that encodes a functionally complete human CH or CL immunoglobulin sequence, with appropriate restriction sites engineered so that any VH or VL sequence can be easily inserted and expressed. In such vectors, splicing usually occurs between the splice donor site in the inserted J region and the splice acceptor site preceding the human C region, and also at the splice regions that occur within the human CH exons. Polyadenylation and transcription termination occur at native chromosomal sites downstream of the coding regions. The resulting chimeric antibody may be joined to any strong promoter, including retroviral ITRs, e.g. SV-40 early promoter, (Okayama *et al.* (1983) *Mol. Cell. Bio.* 3:280), Rous sarcoma virus ITR (Gorman *et al.* (1982) *P.N.A.S.* 79:6777), and moloney murine leukemia virus ITR (Grosschedl *et al.* (1985) *Cell* 41:885); native Ig promoters, etc.

In yet other embodiments of the invention, the active agent is an agent that modulates, and generally decreases or down regulates, the expression of the gene encoding the target protein in the host. For example, antisense molecules can be used to down-regulate expression of MPTS in cells. The anti-sense reagent may be antisense

or a short double-stranded nucleic acid sequence complementary to the mRNA of the targeted gene, and inhibits expression of the targeted gene products. Antisense

molecules inhibit gene expression through various mechanisms, e.g. by reducing the amount of mRNA available for translation, through activation of RNase H, or steric hindrance. One or a combination of antisense molecules may be administered, where a combination may comprise multiple different sequences

5

Antisense molecules may be produced by expression of all or a part of the target gene sequence in an appropriate vector, where the transcriptional initiation is oriented such that an antisense strand is produced as an RNA molecule. Alternatively, the antisense molecule is a synthetic oligonucleotide. Antisense oligonucleotides will  
10 generally be at least about 7, usually at least about 12, more usually at least about 20 nucleotides in length, and not more than about 500, usually not more than about 50, more usually not more than about 35 nucleotides in length, where the length is governed by efficiency of inhibition, specificity, including absence of cross-reactivity, and the like. It has been found that short oligonucleotides, of from 7 to 8 bases in  
15 length, can be strong and selective inhibitors of gene expression (see Wagner *et al.* (1996), *Nature Biotechnol.* 14:840-844).

A specific region or regions of the endogenous sense strand mRNA sequence is chosen to be complemented by the antisense sequence. Selection of a specific sequence  
20 for the oligonucleotide may use an empirical method, where several candidate sequences are assayed for inhibition of expression of the target gene in an *in vitro* or animal model. A combination of sequences may also be used, where several regions of the mRNA sequence are selected for antisense complementation.

25 Antisense oligonucleotides may be chemically synthesized by methods known in the art (see Wagner *et al.* (1993), *supra*, and Milligan *et al.*, *supra*.) Preferred oligonucleotides are chemically modified from the native phosphodiester structure, in order to increase their intracellular stability and binding affinity. A number of such modifications have been described in the literature, which alter the chemistry of the  
30 backbone, sugars or heterocyclic bases.

Among useful changes in the backbone chemistry are phosphorothioates; phosphorodithioates, where both of the non-bridging oxygens are substituted with

sulfur; phosphoroamidites; alkyl phosphotriesters and boranophosphates. Achiral phosphate derivatives include 3'-O'-5'-S-phosphorothioate, 3'-S-5'-O-phosphorothioate, 3'-CH<sub>2</sub>-5'-O-phosphonate and 3'-NH-5'-O-phosphoroamidate. Peptide nucleic acids replace the entire ribose phosphodiester backbone with a peptide linkage. Sugar modifications are also used to enhance stability and affinity. The  $\alpha$ -anomer of deoxyribose may be used, where the base is inverted with respect to the natural  $\beta$ -anomer. The 2'-OH of the ribose sugar may be altered to form 2'-O-methyl or 2'-O-allyl sugars, which provides resistance to degradation without comprising affinity. Modification of the heterocyclic bases must maintain proper base pairing. Some useful substitutions include deoxyuridine for deoxythymidine; 5-methyl-2'-deoxycytidine and 5-bromo-2'-deoxycytidine for deoxycytidine. 5-propynyl-2'-deoxyuridine and 5-propynyl-2'-deoxycytidine have been shown to increase affinity and biological activity when substituted for deoxythymidine and deoxycytidine, respectively.

As an alternative to anti-sense inhibitors, catalytic nucleic acid compounds, e.g. ribozymes, anti-sense conjugates, *etc.* may be used to inhibit gene expression. Ribozymes may be synthesized *in vitro* and administered to the patient, or may be encoded on an expression vector, from which the ribozyme is synthesized in the targeted cell (for example, see International patent application WO 9523225, and Beigelman *et al.* (1995), *Nucl. Acids Res.* 23:4434-42). Examples of oligonucleotides with catalytic activity are described in WO 9506764. Conjugates of anti-sense ODN with a metal complex, e.g. terpyridylCu(II), capable of mediating mRNA hydrolysis are described in Bashkin *et al.* (1995), *Appl. Biochem. Biotechnol.* 54:43-56.

It is herefore an object of the present invention to provide a method of modulating MPTS activity in a host, said method comprising: administering an effective amount of an MPTS modulatory agent to said host, or more specifically such a method wherein said modulatory agent is a small molecule, an antibody, or a nucleic acid.

As mentioned above, an effective amount of the active agent is administered to

e.g. aggrecanase, activity, as measured by aggrecan cleavage product production, as compared to a control.

In the subject methods, the active agent(s) may be administered to the host  
5 using any convenient means capable of resulting in the desired modulation of MPTS  
activity, e.g. desired reduction in aggrecan cleavage product production. Thus, the agent  
can be incorporated into a variety of formulations for therapeutic administration. More  
particularly, the agents of the present invention can be formulated into pharmaceutical  
compositions by combination with appropriate, pharmaceutically acceptable carriers or  
10 diluents, and may be formulated into preparations in solid, semi-solid, liquid or gaseous  
forms, such as tablets, capsules, powders, granules, ointments, solutions, suppositories,  
injections, inhalants and aerosols.

As such, administration of the agents can be achieved in various ways, including  
15 oral, buccal, rectal, parenteral, intraperitoneal, intradermal, transdermal, intracheal, etc.,  
administration.

In pharmaceutical dosage forms, the agents may be administered in the form of  
their pharmaceutically acceptable salts, or they may also be used alone or in appropriate  
20 association, as well as in combination, with other pharmaceutically active compounds.

For oral preparations, the agents can be used alone or in combination with  
appropriate additives to make tablets, powders, granules or capsules, for example, with  
conventional additives, such as lactose, mannitol, corn starch or potato starch; with  
25 binders, such as crystalline cellulose, cellulose derivatives, acacia, corn starch or gelatins;  
with disintegrators, such as corn starch, potato starch or sodium  
carboxymethylcellulose; with lubricants, such as talc or magnesium stearate; and if  
desired, with diluents, buffering agents, moistening agents, preservatives and flavoring  
agents.

30

The agents can be formulated into preparations for injection by dissolving,  
suspending or emulsifying them in an aqueous or nonaqueous solvent, such as vegetable  
or other similar oils, synthetic aliphatic acid glycerides, esters of higher aliphatic acids or

propylene glycol; and if desired, with conventional additives such as solubilizers, isotonic agents, suspending agents, emulsifying agents, stabilizers and preservatives.

The agents can be utilized in aerosol formulation to be administered via  
5 inhalation. The compounds of the present invention can be formulated into pressurized acceptable propellants such as dichlorodifluoromethane, propane, nitrogen and the like.

Furthermore, the agents can be made into suppositories by mixing with a variety of bases such as emulsifying bases or water-soluble bases. The compounds of the present  
10 invention can be administered rectally via a suppository. The suppository can include vehicles such as cocoa butter, carbowaxes and polyethylene glycols, which melt at body temperature, yet are solidified at room temperature.

Unit dosage forms for oral or rectal administration such as syrups, elixirs, and  
15 suspensions may be provided wherein each dosage unit, for example, teaspoonful, tablespoonful, tablet or suppository, contains a predetermined amount of the composition containing one or more inhibitors. Similarly, unit dosage forms for injection or intravenous administration may comprise the inhibitor(s) in a composition as a solution in sterile water, normal saline or another pharmaceutically acceptable  
20 carrier.

The term "unit dosage form," as used herein, refers to physically discrete units suitable as unitary dosages for human and animal subjects, each unit containing a predetermined quantity of compounds of the present invention calculated in an amount  
25 sufficient to produce the desired effect in association with a pharmaceutically acceptable diluent, carrier or vehicle. The specifications for the novel unit dosage forms of the present invention depend on the particular compound employed and the effect to be achieved, and the pharmacodynamics associated with each compound in the host.

The compounds of the present invention, and the various pharmaceutical compositions, agents, stabilizers, wetting agents and the like, are readily available to the public.

Where the agent is a polypeptide, polynucleotide, analog or mimetic thereof, e.g. antisense composition, it may be introduced into tissues or host cells by any number of routes, including viral infection, microinjection, or fusion of vesicles. Jet injection may also be used for intramuscular administration, as described by Furth *et al.* (1992), *Anal*  
 5 *Biochem* 205:365-368. The DNA may be coated onto gold microparticles, and delivered intradermally by a particle bombardment device, or "gene gun" as described in the literature (see, for example, Tang *et al.* (1992), *Nature* 356:152-154), where gold microprojectiles are coated with the therapeutic DNA, then bombarded into skin cells.

10 Those of skill in the art will readily appreciate that dose levels can vary as a function of the specific compound, the severity of the symptoms and the susceptibility of the subject to side effects. Preferred dosages for a given compound are readily determinable by those of skill in the art by a variety of means.

15 The subject methods find use in the treatment of a variety of different disease conditions involving MPTS activity, including disease conditions involving aggrecanase activity. Of particular interest is the use of the subject methods to treat disease conditions characterized by the presence of aggrecan cleavage products, particularly 60 kDa aggrecan cleavage products having an ARGS N-terminus. Specific diseases that are  
 20 characterized by the presence of such methods include: rheumatoid arthritis, osteoarthritis, infectious arthritis, gouty arthritis, psoriatic arthritis, spondylitis, sports injury, joint trauma, pulmonary disease, fibrosis, and the like.

By treatment is meant at least an amelioration of the symptoms associated with  
 25 the pathological condition afflicting the host, where amelioration is used in a broad sense to refer to at least a reduction in the magnitude of a parameter, e.g. symptom, associated with the pathological condition being treated, such as hyperphosphatemia. As such, treatment also includes situations where the pathological condition, or at least symptoms associated therewith, are completely inhibited, e.g. prevented from  
 30 happening, or stopped, e.g. terminated, such that the host no longer suffers from the pathological condition, or at least the symptoms that characterize the pathological condition.

A variety of hosts are treatable according to the subject methods. Generally such hosts are "mammals" or "mammalian," where these terms are used broadly to describe organisms which are within the class mammalia, including the orders carnivore (*e.g.*, dogs and cats), rodentia (*e.g.*, mice, guinea pigs, and rats), and primates (*e.g.*, humans, chimpanzees, and monkeys). In many embodiments, the hosts will be humans.

Kits with unit doses of the active agent, usually in oral or injectable doses, are provided. In such kits, in addition to the containers containing the unit doses will be an informational package insert describing the use and attendant benefits of the drugs in treating pathological condition of interest. Preferred compounds and unit doses are those described herein above.

Finally it is an object of the present invention:

- 15 (i) A method of screening to identify MPTS modulatory agents, said method comprising:  
contacting an MPTS proteins as defined herein with a substrate in the  
presence of an potential modulatory agents; and  
determining the effect of said modulatory agent on the activity of said protein.
- 20 (ii) The method as defined in (i), wherein said substrate comprises a glu-ala bond.
- (iii) The method as defined in (i), wherein said substrate is aggrecan or a  
25 fragment thereof.

Furthermore, a method of treating a host suffering from a disease condition associated with MPTS activity specifically wherein said disease condition is characterized by the presence of aggrecan cleavage products, said method comprising:

*administering to said host an effective amount of a compound of the present invention;*

*wherein said compound is a MPTS modulatory agent;*  
a medicament for the treatment of a disease condition associated with MPTS activity.

specifically wherein said disease condition is characterized by the presence of aggrecan cleavage products, like arthritis.

### Examples

5

#### Example 1

A nucleic acid array carrying 699 known metalloproteinase genes and novel ESTs available in public and proprietary databases was designed. These sequences on the array were selected by a search with a seed set of known metalloprotease protein  
10 sequences from all species. These protein sequences were used to find matching sequences in human nucleotide at the protein (codon) level. Redundant sequences were eliminated, remaining sequences assembled and clustered, and the unique set of 699 sequences were arrayed.

15

The resultant array was used to screen genes expressed in primary cultures of chondrocytes. A fair number of metalloproteinases known to be expressed by these cells were identified. However, a number of ESTs for novel proteins were also identified. Using these ESTs in subsequent database mining and PCR protocols, four different human MPTS proteins were identified, i.e. MPTS15, MPTS10, MPTS19 and MPTS20.

20

#### Example 2

##### Expression of MPTS-10

25

An example of a system for expression of mpts-10 is the COS-7 mammalian cell system. The nucleotide sequence that encodes mpts-10, including the secretion signal sequence, was ligated into a pcDNA3.1 plasmid (In Vitrogen, Carlsbad, CA, USA). Two micrograms of the resulting plasmid was combined with lipofectamine (Life Technologies, Rockville, MD, USA). The mixture was then added to COS-7 cells, which  
30 were grown in 6 well plates to a density of approximately 90% confluency. After 6 hours, fresh medium was added to the cells and after 24 hours the cells were washed and fresh serum free medium containing bovine aggrecan (0.1mg/ml, Sigma, St. Louis, MO, USA) was added. The cells were incubated for an additional 48 hours. Five hundred



microliters of culture fluid from each well was collected and concentrated ten fold. Two microliters of chondroitinase ABC and keratinase (10 u/ml, Sigma, St. Louis, MO, USA) was then added and the samples incubated overnight at 37C. The samples were then boiled in SDS-PAGE sample loading buffer, electrophoresed on a polyacryamide gel and transferred to a PVDF membrane. A Western blot using an antiserum against a neopeptide generated when aggrecanase cleaves aggrecan was then performed.

Another example of a system for expression of mpts-10 was the baculovirus expression system. The DNA sequence that contained the coding sequence for mpts-10 (including the sequences that code for the secretion signal sequence) and that had been cloned in the pcDNA3.1 vector was modified by PCR so that the coding sequence and the translational stop codon were flanked by the Not I (N-terminal side) and Sfi-I (C-terminal side). The primer used for the N-terminal end was GATCGCGGCCGCTATGGTGGACACGTGGCCTCTATGGCTCC and the primer for the C-terminal end was TGAGGCCTTCAGGGCCGATCACTGTGCAGAGCACTCACCCCAT. After amplification using standard PCR methods, the fragment was digested with Not I and Sfi-I. The digested fragment was ligated into a vector pVL1392-U, which had also been digested with NotI and Sfi-I. PVL1392-U is a derivation of the baculovirus transfer plasmid, pVL1392 (PharMingen, San Diego, CA USA) in which the multiple cloning site has been modified to contain Not-I and Sfi-I. The overhangs generated by digestion with Not-I and Sfi-I were complementary to the overhangs generated in the Not I and Sfi I digested PCR amplified DNA. The ligated DNA was transformed into bacterial cells and a clone was selected that contained the plasmid and the correct mpts-10 sequence. This plasmid was produced and purified. The mpts-10 sequence was transferred into a baculovirus vector using standard techniques (*Baculovirus Expression Vectors: A Laboratory Manual* by David O'Reilly, Lois Miller, and Verne Luckow, W.H. Freeman and Co., New York, USA). Five plaque purified virus preparations were produced from the virus preparation. Sf9 insect cells growing in suspension were infected with each of the plaque purified virus preparations at a multiplicity of 0.5

chondroitinase ABC and keratinase (10u/ml) at 37C overnight. The samples were then examined by Western blotting using an antiserum that reacts with a neoepitope generated when aggrecan is cleaved by aggrecanase.

5           Another method for expression of mpts-10 was the drosophila expression system. The DNA fragment containing the sequences encoding mpts-10 and flanked by Not-1 and Sfi-1 that had been generated by PCR (see above) was cloned into plasmid Cmk 33. Cmk33 is a plasmid derived from pMK33/pMtHy (Li, Bin et al Biochem J (1996) 313, 57-64) so that Not-1 and Sfi-1 were in the cloning site. The overhangs  
10   generated by digestion of this plasmid are compatible with the overhangs generated in the digested DNA containing the mpts-10 fragment (see above). A plasmid containing the correct sequence of mpts-10 was amplified and purified. Drosophila (S2) cells were transformed with the plasmid using standard techniques (Li, Bin et al Biochem J (1996) 313, 57-64). Culture fluid was collected 2 days after transfection. These samples were  
15   assayed for aggrecanase activity by incubating with bovine aggrecan (Sigma, St. Louis, MO, USA) at a concentration of 0.1 mg/ml. The samples were then incubated with both chondroitinase ABC and keratinase (10u/ml) at 37C overnight. The samples were then examined by Western blotting using an antiserum that reacts with a neoepitope generated when aggrecan is cleaved by aggrecanase.

20

### Example 3

#### Expression of MPTS-15

25           An example of a system for expression of mpts-15 is the COS-7 mammalian cell system. The nucleotide sequence that encodes mpts-15, including the secretion signal sequence, was ligated into a pcDNA3.1 plasmid (In Vitrogen, Carlsbad, CA, USA). Two micrograms of the resulting plasmid was combined with lipofectamine (Life Technologies, Rockville, MD, USA). The mixture was then added to COS-7 cells, which  
30   were grown in 6 well plates to a density of approximately 90% confluency. After 6 hours, fresh medium was added to the cells and after 24 hours the cells were washed and fresh serum free medium containing bovine aggrecan (0.1mg/ml, Sigma, St. Louis, MO, USA) was added. The cells incubated for an additional 48 hours. Five hundred

microliters of culture fluid from each well was collected and concentrated ten fold. Two microliters of chondroitinase ABC and keratinase (10 u/ml, Sigma, St. Louis, MO, USA) was then added and the samples incubated overnight at 37C. The samples were then boiled in SDS-PAGE sample loading buffer, electrophoresed on a polyacryamide gel and transferred to a PVDF membrane. A Western blot using an antiserum against a neoepitope generated when aggrecanase cleaves aggrecan was then performed.

Another example of a system for expression of mpts-15 was the baculovirus expression system. The DNA sequence that contained the coding sequence for mpts-15 (including the sequences that code for the secretion signal sequence) and that had been cloned in the pcDNA3.1 vector was modified by PCR so that the coding sequence and the translational stop codon were flanked by the Not I (N-terminal side) and Sfi-1 (C-terminal side). The primer used for the N-terminal end was GATCGCGGCCGCTATGGAAATTTGTGGAAGACGTTG and the primer for the C-terminal end was TGAGGCCTTCAGGGCCGATCTTAAAGCAAAGTTTCTTTTGGT. After amplification using standard PCR methods, the fragment was digested with Not I and Sfi-1. The digested fragment was ligated into a vector pVL1392-U, which had also been digested with NotI and Sfi-1. PVL1392-U is a derivation of the baculovirus transfer plasmid, pVL1392 (PharMingen, San Diego, CA, USA) in which the multiple cloning site has been modified to contain Not-1 and Sfi-1. The overhangs generated by digestion with Not-1 and Sfi-1 were complementary to the overhangs generated in the Not I and Sfi I digested PCR amplified DNA. The ligated DNA was transformed into bacterial cells and a clone was selected that contained the plasmid and the correct mpts-15 sequence. This plasmid was produced and purified. The mpts-15 sequence was transferred into a baculovirus vector using standard techniques (*Baculovirus Expression Vectors: A Laboratory Manual* by David O'Reilly, Lois Miller, and Verne Luckow, W.H. Freeman and Co., New York, USA). Five plaque purified virus preparations were produced from the virus preparation. Sf9 insect cells growing in suspension were infected with each of the plaque purified virus preparations at a multiplicity of 0.5. Culture fluid was harvested at 3 days after infection. These samples were assayed for

examined by Western blotting using an antiserum that reacts with a neoepitope generated when aggrecan is cleaved by aggrecanase.

Another method for expression of mpts-15 was the drosophila expression system. The DNA fragment containing the sequences encoding mpts-15 and flanked by Not-1 and Sfi-1 that had been generated by PCR (see above) was cloned into plasmid Cmk 33. Cmk33 is a plasmid derived from pMK33/pMtHy (Li, Bin et al Biochem J (1996) 313, 57-64) so that Not-1 and Sfi-1 were in the cloning site. The overhangs generated by digestion of this plasmid are compatible with the overhangs generated in the Not 1 and Sfi 1 digested DNA containing the mpts-15 fragment (see above). A plasmid containing the correct sequence of mpts-15 was amplified and purified. Drosophila (S2) cells were transformed with the plasmid using standard techniques (Li, Bin et al Biochem J (1996) 313, 57-64). Culture fluid was collected 2 days after transfection. These samples were assayed for aggrecanase activity by incubating with bovine aggrecan (Sigma, St. Louis, MD, USA) at a concentration of 0.1 mg/ml. The samples were then incubated with both chondroitinase ABC and keratinase (10u/ml) at 37C overnight. The samples were then examined by Western blotting using an antiserum that reacts with a neoepitope generated when aggrecan is cleaved by aggrecanase.

#### Example 4

##### Expression of MPTS-19

An example of a system for expression of mpts-19 is the COS-7 mammalian cell system. The nucleotide sequence that encodes mpts-19, including the secretion signal sequence and the C-terminal stop codon, was ligated into a pcDNA3.1 plasmid (In Vitrogen, Carlsbad, CA, USA). Two micrograms of the resulting plasmid was combined with lipofectamine (Life Technologies, Rockville, MD, USA). The mixture was then added to COS-7 cells, which were grown in 6 well plates to a density of approximately 90% confluency. After 6 hours, fresh medium was added to the cells and after 24 hours the cells were washed and fresh serum free medium containing bovine aggrecan (0.1mg/ml, Sigma, St. Louis, MO, USA) was added. The cells incubated for an

additional 48 hours. Five hundred microliters of culture fluid from each well was collected and concentrated ten fold. Two microliters of chondroitinase ABC and keratinase (10 u/ml, Sigma, St. Louis, MO, USA) was then added and the samples incubated overnight at 37C. The samples were then boiled in SDS-PAGE sample  
 5 loading buffer, electrophoresed on a polyacrylamide gel and transferred to a PVDF membrane. A Western blot using an antiserum against a neoepitope generated when aggrecanase cleaves aggrecan was then performed.

Another example of a system for expression of mpts-19 was the baculovirus  
 10 expression system. The DNA sequence that contained the coding sequence for mpts-19 (including the sequences that code for the secretion signal sequence) and that had been cloned in the pcDNA3.1 vector was modified by PCR so that the coding sequence and the translational stop codon were flanked by the Not I (N-terminal side) and Sfi-I (C-terminal side). The primer used for the N-terminal end was  
 15 GATCGCGGCCGCTATGCCCGGCGGCCCCAGTCCCCG and the primer for the C-terminal end was  
TGAGGCCTTCAGGGCCGATCTCAGCGGCGGGCAACCCGCTG. After amplification using standard PCR methods, the fragment was digested with Not I and Sfi-I. The digested fragment was ligated into a vector pVL1392-U, which had also been  
 20 digested with NotI and Sfi-I. PVL1392-U is a derivation of the baculovirus transfer plasmid, pVL1392 (PharMingen, San Diego, CA USA) in which the multiple cloning site has been modified to contain Not-I and Sfi-I. The overhangs generated by digestion with Not-I and Sfi-I were complementary to the overhangs generated in the Not I and Sfi I digested PCR amplified DNA. The ligated DNA was transformed into bacterial  
 25 cells and a clone was selected that contained the plasmid and the correct mpts-19 sequence. This plasmid was produced and purified. The mpts-19 sequence was transferred into a baculovirus vector using standard techniques (*Baculovirus Expression Vectors: A Laboratory Manual* by David O'Reilly, Lois Miller, and Verne Luckow, W.H. Freeman and Co., New York, USA). Five plaque purified virus preparations were  
 30 produced from the virus preparation. Sf9 insect cells growing in suspension were

Aggrecanase (A1111) was obtained from Calbiochem, La Jolla, CA, USA at

a concentration of 0.1 mg/ml. The samples were then incubated with both chondroitinase ABC and keratinase (10u/ml) at 37C overnight. The samples were then examined by Western blotting using an antiserum that reacts with a neoepitope generated when aggrecan is cleaved by aggrecanase.

5

Another method for expression of mpts-19 was the drosophila expression system. The DNA fragment containing the sequences encoding mpts-19 and flanked by Not-1 and Sfi-1 that had been generated by PCR (see above) was cloned into plasmid Cmk 33. Cmk33 is a plasmid derived from pMK33/pMtHy (Li, Bin et al Biochem J (1996) 313, 57-64) so that Not-1 and Sfi-1 were in the cloning site. The overhangs generated by digestion of this plasmid with Not 1 and Sfi 1 are compatible with the overhangs generated in the digested DNA containing the mpts-19 fragment (see above). A plasmid containing the correct sequence of mpts-19 was amplified and purified. Drosophila (S2) cells were transformed with the plasmid using standard techniques (Li, Bin et al Biochem J (1996) 313, 57-64). Culture fluid was collected 2 days after transfection. These samples were assayed for aggrecanase activity by incubating with bovine aggrecan (Sigma, St. Louis, MO, USA) at a concentration of 0.1 mg/ml. The samples were then incubated with both chondroitinase ABC and keratinase (10u/ml) at 37C overnight. The samples were then examined by Western blotting using an antiserum that reacts with a neoepitope generated when aggrecan is cleaved by aggrecanase.

15

20

#### Example 5

##### 25 Expression of MPTS-20

An example of a system for expression of mpts-20 is the COS-7 mammalian cell system. The nucleotide sequence that encodes mpts-10, including the secretion signal sequence and the C-terminal stop codon, was ligated into a pcDNA3.1 plasmid (In Vitrogen, Carlsbad, CA, USA). Two micrograms of the resulting plasmid was combined with lipofectamine (Life Technologies, Rockville, MD, USA). The mixture was then added to COS-7 cells, which were grown in 6 well plates to a density of approximately 90% confluency. After 6 hours, fresh medium was added to the cells and

30

after 24 hours the cells were washed and fresh serum free medium containing bovine aggrecan (0.1mg/ml, Sigma, St. Louis, MO, USA) was added. The cells incubated for an additional 48 hours. Five hundred microliters of culture fluid from each well was collected and concentrated ten fold. Two microliters of chondroitinase ABC and  
 5 keratinase (10 u/ml, Sigma, St. Louis, MO, USA) was then added and the samples incubated overnight at 37C. The samples were then boiled in SDS-PAGE sample loading buffer, electrophoresed on a polyacryamide gel and transferred to a PVDF membrane. A Western blot using an antiserum against a neoepitope generated when aggrecanase cleaves aggrecan was then performed.

10

Another example of a system for expression of mpts-20 was the baculovirus expression system. The DNA sequence that contained the coding sequence for mpts-20 (including the sequences that code for the secretion signal sequence) and that had been cloned in the pcDNA3.1 vector was modified by PCR so that the coding sequence and  
 15 the translational stop codon were flanked by the Not I (N-terminal side) and Sfi-I (C-terminal side). The primer used for the N-terminal end was  
GATCGCGGCCGCTGCCGCTGTGATGAGTGTGCCTG and the primer for the C-terminal end was

TGAGGCCITTCAGGGCCGATCTTATAAAGGCCITGAGAAAACAG. After  
 20 amplification using standard PCR methods, the fragment was digested with Not I and Sfi-I. The digested fragment was ligated into a vector pVL1392-U, which had also been digested with NotI and Sfi-I. PVL1392-U is a derivation of the baculovirus transfer plasmid, pVL1392 (PharMingen, San Diego, CA USA) in which the multiple cloning site has been modified to contain Not I and Sfi-I. The overhangs generated by digestion  
 25 with Not I and Sfi-I were complementary to the overhangs generated in the Not I and Sfi I digested PCR amplified DNA. The ligated DNA was transformed into bacterial cells and a clone was selected that contained the plasmid and the correct mpts-20 sequence. This plasmid was produced and purified. The mpts-20 sequence was transferred into a baculovirus vector using standard techniques (*Baculovirus Expression*  
 3 *Vectors: A Laboratory Manual* (Dowd, D. and D. L. Kings, Eds., Academic Press, 1998).

mixed with each of the plaque-purified virus preparations at a multiplicity of 0.5.

Culture fluid was harvest 3 days after infection. These samples were assayed for aggrecanase activity by incubating with bovine aggrecan (Sigma, St. Louis, MO, USA) at a concentration of 0.1 mg/ml. The samples were then incubated with both chondroitinase ABC and keratinase (10u/ml) at 37C overnight. The samples were then  
5 examined by Western blotting using an antiserum that reacts with a neoepitope generated when aggrecan is cleaved by aggrecanase.

Another method for expression of mpts-20 was the drosophila expression system. The DNA fragment containing the sequences encoding mpts-20 and flanked by  
10 Not-1 and Sfi-1 that had been generated by PCR (see above) was cloned into plasmid Cmk 33. Cmk33 is a plasmid derived from pMK33/pMtHy (Li, Bin et al Biochem J (1996) 313, 57-64) so that Not-1 and Sfi-1 were in the cloning site. The overhangs generated by digestion of this plasmid with Not 1 and Sfi 1 are compatible with the overhangs generated in the digested DNA containing the mpts-20 fragment (see above).  
15 A plasmid containing the correct sequence of mpts-20 was amplified and purified. Drosophila (S2) cells were transformed with the plasmid using standard techniques (Li, Bin et al Biochem J (1996) 313, 57-64). Culture fluid was collected 2 days after transfection. These samples were assayed for aggrecanase activity by incubating with bovine aggrecan (Sigma, St. Louis, MO, USA) at a concentration of 0.1 mg/ml. The  
20 samples were then incubated with both chondroitinase ABC and keratinase (10u/ml) at 37C overnight. The samples were then examined by Western blotting using an antiserum that reacts with a neoepitope generated when aggrecan is cleaved by aggrecanase.

#### 25 Example 6

##### Purification of mpts-10, 15, 19 and 20:

Mpts-10, 15, 19, and 20 were purified from the culture fluid of the expression systems  
30 described above using chromatographic procedures. For example, the culture fluid was adjusted with regard to pH, filtered and then loaded onto a column packed with sulfoethyl sepharose FF (Amersham-Pharmacia Biotech, Piscataway, NJ, USA). After washing with a buffer consisting of 10 mM CaCl<sub>2</sub>, 0.1 M NaCl, and 0.05% Brij35 at a pH



which results in retention of the mpts's on the column, the mpts's were eluted with a 0.1 M to 1.0 M NaCl gradient. Fractions from the column were assayed for the presence of aggrecanase activity as described above and noted. For the purification a column packed with phenylsepharose or sephacryl S-200 can also be used.

## SEQUENCE LISTING

<110> Hoffmann-La Roche AG

5 <120> Novel Metalloproteases Having  
Thrombospondin Domains and Nucleic Acid Compositions  
Encoding the Same

<130> 20594

10

<150> 50/184,152

<151> 2000-02-18

<160> 10

15

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 959

20 <212> PRT

<213> human

<220>

<225> Xaa at 909 is any amino acid

25

<223> X at 211 is any amino acid

<223> X at 218 is any amino acid

30 <223> Xaa is any amino acid

&lt;400&gt; 1

Met Glu Ile Leu Trp Lys Thr Leu Thr Trp Ile Leu Ser Leu Ile Met  
 1 5 10 15  
 Ala Ser Ser Glu Phe His Ser Asp His Arg Leu Ser Tyr Ser Ser Gln  
 5 20 25 30  
 Glu Glu Phe Leu Thr Tyr Leu Glu His Tyr Gln Leu Thr Ile Pro Ile  
 35 40 45  
 Arg Val Asp Gln Asn Gly Ala Phe Leu Ser Phe Thr Val Lys Asn Asp  
 50 55 60  
 10 Lys His Ser Arg Arg Arg Arg Ser Met Asp Pro Ile Asn Pro Gln Gln  
 65 70 75 80  
 Ala Val Ser Lys Leu Phe Phe Lys Leu Ser Ala Tyr Gly Lys His Phe  
 85 90 95  
 His Leu Asn Leu Thr Leu Asn Thr Asp Phe Val Ser Lys His Phe Thr  
 15 100 105 110  
 Val Glu Tyr Trp Gly Lys Asp Gly Pro Gln Trp Lys His Asp Phe Leu  
 115 120 125  
 Asp Asn Cys His Tyr Thr Gly Tyr Leu Gln Asp Gln Arg Ser Thr Thr  
 130 135 140  
 20 Lys Val Ala Leu Ser Asn Cys Val Gly Leu His Gly Val Ile Ala Thr  
 145 150 155 160  
 Glu Asp Glu Glu Tyr Phe Ile Gln Pro Leu Lys Asn Thr Thr Glu Asp  
 165 170 175  
 Ser Lys His Phe Ser Tyr Gln Asn Gly His Pro His Val Ile Tyr Lys  
 25 180 185 190  
 Lys Ser Ala Leu Gln Gln Arg His Leu Tyr Asp His Ser His Cys Gly  
 195 200 205  
 Val Thr Arg Leu Thr Arg Ser Gly Lys Ile Thr Thr Leu Asn Asp Thr  
 210 215 220  
 30 Leu Thr Val Ser Lys Ser Leu Thr Thr Asn Asn Thr His Ile His His  
 225 230 235 240  
 Arg Gln Lys Arg Ser Val Ser Ile Gln Arg Phe Val Gln Thr Leu Val  
 245 250 255  
 Val Ala Asp Lys Met Met Val Gly Tyr His Gly Arg Lys Asp Ile Gln  
 35 260 265 270  
 His Tyr Thr Leu Ser Thr Met Asn Thr Val Ala Thr Tyr Thr Thr Thr

Leu Thr Glu Asp Gln Pro Asn Leu Glu Ile Asn His His Ala Asp Lys  
 305 310 315 320  
 Ser Leu Asp Ser Phe Cys Lys Trp Gln Lys Ser Ile Leu Ser His Gln  
 325 330 335  
 5 Ser Asp Gly Asn Thr Ile Pro Glu Asn Gly Ile Ala His His Asp Asn  
 340 345 350  
 Ala Val Leu Ile Thr Arg Tyr Asp Ile Cys Thr Tyr Lys Asn Lys Pro  
 355 360 365  
 Cys Gly Thr Leu Gly Leu Ala Ser Val Ala Gly Met Cys Glu Pro Glu  
 10 370 375 380  
 Arg Ser Cys Ser Ile Asn Glu Asp Ile Gly Leu Gly Ser Ala Phe Thr  
 385 390 395 400  
 Ile Ala His Glu Ile Gly His Asn Phe Gly Met Asn His Asp Gly Ile  
 405 410 415  
 15 Gly Asn Ser Cys Gly Thr Lys Gly His Glu Ala Ala Lys Leu Met Ala  
 420 425 430  
 Ala His Ile Thr Ala Asn Thr Asn Pro Phe Ser Trp Ser Ala Cys Ser  
 435 440 445  
 Arg Asp Tyr Ile Thr Ser Phe Leu Asp Ser Gly Arg Gly Thr Cys Leu  
 20 450 455 460  
 Asp Asn Glu Pro Pro Lys Arg Asp Phe Leu Tyr Pro Ala Val Ala Pro  
 465 470 475 480  
 Gly Gln Val Tyr Asp Ala Asp Glu Gln Cys Arg Phe Gln Tyr Gly Ala  
 485 490 495  
 25 Thr Ser Arg Gln Cys Lys Tyr Gly Glu Val Cys Arg Glu Leu Trp Cys  
 500 505 510  
 Leu Ser Lys Ser Asn Arg Cys Val Thr Asn Ser Ile Pro Ala Ala Glu  
 515 520 525  
 Gly Thr Leu Cys Gln Thr Gly Asn Ile Glu Lys Gly Trp Cys Tyr Gln  
 30 530 535 540  
 Gly Asp Cys Val Pro Phe Gly Thr Trp Pro Gln Ser Ile Asp Gly Gly  
 545 550 555 560  
 Trp Gly Pro Trp Ser Leu Trp Gly Glu Cys Ser Arg Thr Cys Gly Gly  
 565 570 575  
 35 Gly Val Ser Ser Ser Leu Arg His Cys Asp Ser Pro Ala Pro Ser Gly  
 580 585 590  
 Gly Gly Lys Tyr Cys Leu Gly Glu Arg Lys Arg Tyr Arg Ser Cys Asn  
 595 600 605  
 Thr Asp Pro Cys Pro Leu Gly Ser Arg Asp Phe Arg Glu Lys Gln Cys  
 40 610 615 620

Ala Asp Phe Asp Asn Met Pro Phe Arg Gly Lys Tyr Tyr Asn Trp Lys  
 625 630 635 640  
 Pro Tyr Thr Gly Gly Gly Val Lys Pro Cys Ala Leu Asn Cys Leu Ala  
 645 650 655  
 5 Glu Gly Tyr Asn Phe Tyr Thr Glu Arg Ala Pro Ala Val Ile Asp Gly  
 660 665 670  
 Thr Gln Cys Asn Ala Asp Ser Leu Asp Ile Cys Ile Asn Gly Glu Cys  
 675 680 685  
 Lys His Val Gly Cys Asp Asn Ile Leu Gly Ser Asp Ala Arg Glu Asp  
 10 690 695 700  
 Arg Cys Arg Val Cys Gly Gly Asp Gly Ser Thr Cys Asp Ala Ile Glu  
 705 710 715 720  
 Gly Phe Phe Asn Asp Ser Leu Pro Arg Gly Gly Tyr Met Glu Val Val  
 725 730 735  
 15 Gln Ile Pro Arg Gly Ser Val His Ile Glu Val Arg Glu Val Ala Met  
 740 745 750  
 Ser Lys Asn Tyr Ile Ala Leu Lys Ser Glu Gly Asp Asp Tyr Tyr Ile  
 755 760 765  
 Asn Gly Ala Trp Thr Ile Asp Trp Pro Arg Lys Phe Asp Val Ala Gly  
 20 770 775 780  
 Thr Ala Phe His Tyr Lys Arg Pro Thr Asp Glu Pro Glu Ser Leu Glu  
 785 790 795 800  
 Ala Leu Gly Pro Thr Ser Glu Asn Leu Ile Val Met Val Leu Leu Glu  
 805 810 815  
 25 Glu Glu Asn Leu Gly Ile Arg Tyr Lys Phe Asn Val Pro Ile Thr Arg  
 820 825 830  
 Thr Gly Ser Gly Asp Asn Glu Val Gly Phe Thr Trp Asn His Gln Ser  
 835 840 845  
 Trp Ser Glu Cys Ser Ala Thr Tyr Ala Gly Gly Lys Met Ile Thr Arg  
 850 855 860  
 Gln Ile Thr Gln Arg Ala Arg Trp Asn Thr Lys His Ile Leu Ser Tyr  
 865 870 875 880  
 Ala Leu Cys Leu Leu Lys Lys Leu Ile Gly Asn Ile Ser Cys Arg Phe  
 885 890 895  
 35 Ala Ser Ser Cys Asn Leu Pro Lys Glu Ala Leu Leu Asn Leu Tyr Tyr  
 900 905 910

Tyr Leu Glu Gly Gly Leu Phe Ala Phe Arg Glu His Ile Leu Gly

945

950

955

<210> 2

5 <211> 2879

<212> DNA

<213> human

<400> 2

```

10 atggaattt tgggaagac gttgacctg atttgagcc tcatcatggc ttcacggaa
   60
   ttccatagtg accacaggt ttcatacagt tctcaagagg aattccatgac tcatcttgaa
   120
   cactaccagc taactattcc aataagggtt gatcaaaatg gagcatttct cagctttact
15   180
   gggaaattg ataaacactc aaggagaaga cggagtatgg accctattga tccacagcag
   240
   gcagatctta agttattttt taaactttca gccatggga agcactttca tctaaacttg
   300
20 actcacaaca cagattttgt gtccaaacat ttacagtag aatattgggg aaagatgga
   360
   ccccatggga aacatgattt tttagacaac tgtcattaca caggatattt gaaagatcaa
   420
   cgtatcttca ctaaagtggc cttaagcaac tgggttgggt tgcattgggt tcttgcctaa
25   480
   aaagatgaaa agtattttat cgaactttta aagaattcca cagagatttt caggtatttt
   540
   agttatgaaa atggccaccc tcatgttatt tacaataagt ctgcctttca acaagacact
   600
30 ctgtatgata actctcattg tggggtttcg gatttcacaa gaagtggcaa acattgggtg
   660
   ctgaatgaca catccactgt ttcttattca ctaccaatta acaacacaca tatccaccac
   720
   agacataaga gatcagttag cattgaacgg ttgtgggaga cattggtagt gccagacaaa
35   780
   atgatggtag gctaccatgg ccgcaaagac attgaacatt acattttgag tctgatgaat
   840
   attgttgcca aactttaccg tgattccagc ctaggaaaac ttgtgaatat tatagtggcc
   900

```

cgcttaattg ttctcacaga agatcagcca aacttggaga taaaccacca tgcagacaag  
 960  
 tccctcgata gcttctgtaa atggcagaaa tccattctct cccaccaaag tgatggaaac  
 1020  
 5 accattccag aaaatgggat tgcccaccac gataatgcag ttcttattac tagatatgat  
 1080  
 atctgcactt ataaaaataa gccctgtgga acactgggct tggcctctgt ggcctggaatg  
 1140  
 tctgagcctg aaaggagctg cagcattaat gaagacattg gccctgggttc agcttttacc  
 1200  
 10 attgcacatg agattggtea caatttttgt atgaaccatg atggaattgg aaattcttgt  
 1260  
 gggacgaaag gtcatgaagc agcaaaactt atggcagctc acattactgc gaataccaat  
 1320  
 15 cctttttcct ggtctgcttg cagtgcagac tacatcacca gctttctaga ttcaggccgt  
 1380  
 ggtacttgcc ttgataatga gccctccaag cgtgaacttc ttatccagc tctggcccca  
 1440  
 ggtcaggtgt atgatgctga tgagcaatgt cgtttccagt atggagcaac ctcccgccaa  
 1500  
 20 tgtaaatatz gggaggtgtg tagagagctc tgggtgctca gcaaaagcaa cggctgtgtc  
 1560  
 accaacagta ttccagcagc tgaggggaca ctggttcaaa ctgggaatat tgaaaaaggg  
 1620  
 25 tgggtgcttc agggagattg tgttctctt ggtaactggt ccagagagcat agatggggcc  
 1680  
 tgggtctctt ggtcactata gggaggtgtg agcaaatc ctgagggagj cctctctga  
 1740  
 tccctacgac atgggagac tccaggaat tctgactc acattattg cctctgagaa  
 1800  
 30 aggaacagat accgctctc taccagagat ccttctctt tgggtctcag agatttttga  
 1860  
 gagaacagct ctgcagactt tgacaatatz cctttccag gaaagtatta taactggaaa  
 1920  
 35 ccttatactg gaggctgggt aaacattgt gatttaact atttggctga aggttataat  
 1980

52

gotaggggaag atagatgtcg agtctgtgga ggggacggaa gcacatgtga tgcattgaa  
 2160  
 ggggtttctca atgattcact gccagggga ggctacatgg aagtgggca gataccaaga  
 2220  
 5 ggtctgttc acattgaagt tagagaagtt gccatgtcaa agaattatat tgccttaaaa  
 2280  
 tctgaaggag atgattacta tattaatggt gcttggacta ttgattggcc taggaaattt  
 2340  
 gatgttgctg ggacagcttt tcattacaag agaccaactg atgaaccaga atccttggaa  
 10 2400  
 gctctaggtc ctacctcaga aaatctcacc gtcattggtt tgttcaaga acagaatttg  
 2460  
 ggaatnaggt ataagttcaa tgttccacc accgaactg gcagtgga taatgaagtt  
 2520  
 15 agcttaccat ggaatcatca gtcttggta gaatgctcag ctcttgtgc tggaggtaag  
 2580  
 atgcccacta ggcagccac ccagagggca agatggagaa caaacaca: tctgagctat  
 2640  
 ccttctggtt tcttaaaaaa gctaattgga aacattctt gcaggttgc tccaagctgt  
 20 2700  
 caattaccaa aagaaactt gcttaatta tattatatt ccttcttcc caacctcctg  
 2760  
 taatttgc agatttctg gtaaaatata tcttggcaca atgagtgct atgcttgc  
 2820  
 25 tctctcaag actatctga aggtgggctg ttgctcttc atgaatata: tcttggat  
 2880

&lt;210&gt; 3

&lt;211&gt; 947

30 &lt;212&gt; PRT

&lt;213&gt; human

&lt;400&gt; 3

Met Ala Pro Ala Cys Gln Ile Leu Arg Trp Ala Leu Ala Leu Gly Leu  
 35 1 5 10 15  
 Gly Leu Met Phe Glu Val Thr His Ala Phe Arg Ser Gln Asp Glu Phe  
 20 25 30  
 Leu Ser Ser Leu Glu Ser Tyr Glu Ile Ala Phe Pro Thr Arg Val Asp  
 35 40 45



53

His Asn Gly Ala Leu Leu Ala Phe Ser Pro Pro Pro Pro Arg Arg Gln  
 50 55 60  
 Arg Arg Gly Thr Gly Ala Thr Ala Glu Ser Arg Leu Phe Tyr Lys Val  
 65 70 75 80  
 5 Ala Ser Pro Ser Thr His Phe Leu Leu Asn Leu Thr Arg Ser Ser Arg  
 85 90 95  
 Leu Leu Ala Gly His Val Ser Val Glu Tyr Trp Thr Arg Glu Gly Leu  
 100 105 110  
 Ala Trp Gln Arg Ala Ala Arg Pro His Cys Leu Tyr Ala Gly His Leu  
 10 115 120 125  
 Gln Gly Gln Ala Ser Ser Ser His Val Ala Ile Ser Thr Cys Gly Gly  
 130 135 140  
 Leu His Gly Leu Ile Val Ala Asp Glu Glu Glu Tyr Leu Ile Glu Pro  
 145 150 155 160  
 15 Leu His Gly Gly Pro Lys Gly Ser Arg Ser Pro Glu Glu Ser Gly Pro  
 165 170 175  
 His Val Val Tyr Lys Arg Ser Ser Leu Arg His Pro His Leu Asp Thr  
 180 185 190  
 Ala Cys Gly Val Arg Asp Glu Lys Pro Trp Lys Gly Arg Pro Trp Trp  
 195 200 205  
 Leu Arg Thr Leu Lys Pro Pro Pro Ala Arg Pro Leu Gly Asn Glu Thr  
 210 215 220  
 Glu Arg Gly Gln Pro Gly Leu Lys Arg Ser Val Ser Arg Glu Arg Tyr  
 225 230 235 240  
 25 Val Glu Thr Leu Val Val Ala Asp Lys Met Met Val Ala Tyr His Gly  
 245 250 255  
 Arg Arg Asp Val Glu His Tyr Val Leu Ala Ile Met Asn Ile Val Ala  
 260 265 270  
 Lys Leu Thr Glu Asp Ser Ser Ser Gly Ser Thr Val Asn Ile Leu Val  
 275 280 285  
 Thr Arg Leu Ile Leu Leu His Gln Asp Gln Pro Thr Leu Glu Ile Thr  
 290 295 300  
 His His Ala Gly Lys Ser Leu Asp Ser Phe Cys Lys Trp Gln Lys Ser  
 305 310 315 320  
 35 Ile Val Asn His Ser Gly His Gly Asn Ala Ile Pro Glu Asn Gly Val  
 325 330 335

Met Cys Glu Arg Glu Arg Ser Cys Ser Val Asn Glu Asp Ile Gly Leu  
 370 375 380  
 Ala Thr Ala Phe Thr Ile Ala His Glu Ile Gly His Thr Phe Gly Met  
 385 390 395 400  
 5 Asn His Asp Gly Val Gly Asn Ser Cys Gly Ala Arg Gly Gln Asp Pro  
 405 410 415  
 Ala Lys Leu Met Ala Ala His Ile Thr Met Lys Thr Asn Pro Phe Val  
 420 425 430  
 Trp Ser Ser Cys Ser Arg Asp Tyr Ile Thr Ser Phe Leu Asp Ser Gly  
 10 435 440 445  
 Leu Gly Leu Cys Leu Asn Asn Arg Pro Pro Arg Gln Asp Phe Val Tyr  
 450 455 460  
 Pro Thr Val Ala Pro Gly Gln Ala Tyr Asp Ala Asp Glu Gln Cys Arg  
 465 470 475 480  
 15 Phe Gln His Gly Val Lys Ser Arg Gly Leu Gln Arg Ala Val Val Ser  
 485 490 495  
 Glu Gln Glu Gln Pro Val His His Gln Gln His Pro Gly Arg Arg Gly  
 500 505 510  
 His Ala Val Pro Asp Ala His His Arg Gln Gly Val Val Leu Gln Thr  
 20 515 520 525  
 Gly Leu Cys Pro Leu Trp Val Ala Pro Arg Gly Cys Gly Arg Ser Leu  
 530 535 540  
 Gly Ala Val Asp Ser Met Gly Asp Cys Ser Arg Thr Cys Gly Gly Gly  
 545 550 555 560  
 25 Val Ser Ser Ser Ser Arg His Cys Asp Ser Pro Arg Pro Thr Ile Gly  
 565 570 575  
 Gly Lys Tyr Cys Leu Gly Gln Arg Arg Arg His Arg Ser Cys Asn Thr  
 580 585 590  
 Asp Asp Cys Pro Pro Gly Ser Gln Asp Phe Arg Glu Val Gln Cys Ser  
 30 595 600 605  
 Glu Phe Asp Ser Ile Pro Phe Arg Gly Lys Phe Tyr Lys Trp Lys Thr  
 610 615 620  
 Tyr Arg Gly Gly Gly Val Lys Ala Cys Ser Leu Thr Cys Leu Ala Glu  
 625 630 635 640  
 35 Gly Phe Asn Phe Tyr Thr Glu Arg Ala Ala Ala Val Val Asp Gly Thr  
 645 650 655  
 Pro Cys Arg Pro Asp Thr Val Asp Ile Cys Val Ser Gly Glu Cys Lys  
 660 665 670  
 His Val Gly Cys Asp Arg Val Leu Gly Ser Asp Leu Arg Glu Asp Lys  
 40 675 680 685

55

Cys Arg Val Cys Gly Gly Asp Gly Ser Ala Cys Glu Thr Ile Glu Gly  
 690 695 700  
 Val Phe Ser Pro Ala Ser Pro Gly Ala Gly Tyr Glu Asp Val Val Trp  
 705 710 715 720  
 5 Ile Pro Lys Gly Ser Val His Ile Phe Ile Gln Asp Leu Asn Leu Ser  
 725 730 735  
 Leu Ser His Leu Ala Leu Lys Gly Asp Gln Glu Ser Leu Leu Leu Glu  
 740 745 750  
 Gly Leu Pro Gly Thr Pro Gln Pro His Arg Leu Pro Leu Ala Gly Thr  
 10 755 760 765  
 Thr Phe Gln Leu Arg Gln Gly Pro Asp Gln Val Gln Ser Leu Glu Ala  
 770 775 780  
 Leu Gly Pro Ile Asn Ala Ser Leu Ile Val Met Val Leu Ala Arg Thr  
 785 790 795 800  
 15 Glu Leu Pro Ala Leu Arg Tyr Arg Phe Asn Ala Pro Ile Ala Arg Asp  
 805 810 815  
 Ser Leu Pro Pro Tyr Ser Trp His Tyr Ala Pro Trp Thr Lys Cys Ser  
 820 825 830  
 Pro Ser Val Gln Ala Val Ala Arg Cys Arg Arg Trp Ser Ala Ala Thr  
 20 835 840 845  
 Lys Leu Asp Ser Ser Ala Val Ala Pro His Tyr Cys Ser Ala His Ser  
 850 855 860  
 Lys Leu Ala Gln Lys Gln Ala Arg Leu Gln His Gly Ala Ile Pro Gln  
 865 870 875 880  
 25 Asp Trp Val Val Gly Thr Val Ala Leu Gln Pro Gln Leu Ala Met Gln  
 885 890 895  
 Gly Val Arg Ser Arg Ser Val Val Cys His Ala Pro Arg Ile Tyr Arg  
 900 905 910  
 His Gln Tyr Ala Leu Arg Arg Ser Ala Tyr Ile Gln Pro Arg Pro Leu  
 915 920 925  
 Val Leu Arg Ile Ala Thr Ala Ile Leu Ala Leu Arg Ser Gly Gly His  
 930 935 940  
 Arg Leu Val  
 945

35

Cys Arg Val

Cys Arg Val

<400> 4  
 ttccatccta atacgactca ctatagggct cgagcggcgc cccgggcagg tgtggacacg  
 60  
 5 tggcctctat ggctcccgcc tgccagatcc tccgctgggc cctcgccctg gggtggggc  
 120  
 tcatgttoga ggtcacgcat gccctccggt ctcaagatga gtccctgtcc agtctggaga  
 180  
 gctatgagat cgccttcccc acccgcggtg accacaaagg ggcactgctg gccttctcgc  
 240  
 10 cactctctcc ccggaggcag cgcgcgggca cggggggcac agccagatcc cgcctcttct  
 300  
 acaaaggggc ctgcgccagc acccacttcc tgcctgaact gacccgtagc tcccgctctc  
 360  
 15 tggcagggca cgtctccgct gagtaactgga caccggaggc cctggcctgg cagagggcgg  
 420  
 cccggtccca ctgcctctac gctggctcacc tgcagggcca ggcagagcgc tcccatgttg  
 480  
 ccatacgcac ctgtggaggc ctgcacggcc tgatcgtggc agacgaggaa gactacctga  
 540  
 20 ttgagctcct gcacggtgga cccaagggtt ctgggagacc ggagyaagt ggaccacatg  
 600  
 tgggtgacaa gggttctctt ctggctcacc ctcaacttga cacagcctgt ggagtgcagc  
 660  
 25 atgagtaacc gtggaaaggc cggccctggc ggttgagjca ctctgaagca cgccttgcca  
 720  
 ggcccctggc caatgaaana gacgctggcc agccaggctt gaagcgatcg ctccagccag  
 780  
 agcgcttccg ggagaccccg gtggtggctg acaagatgat ggtggcctat caccggggcc  
 840  
 30 gggatctgga gcagtatgct ctggccatca tgaacattgt tgcctaaact ttccaggact  
 900  
 cgagtctggg aagcaccggt aacatccctg taactcgctt catctgctc accgaggacc  
 960  
 35 agcccactct ggagatcacc caccatggcg ggaagtccct ggacagcttc tgtaagtggc  
 1020  
 agaaatccat cgtgaaccac agcggccatg gcaatgcat tccagagaa cgtgtggcta  
 1080  
 accatgacac agcagtgtct atcacagct atgacatctg catctacaag aacaaacct  
 1140

ggggcacact aggcctggcc cgggtgggag gaatgtgtga ggcgagaga agctgcagcg  
 1200  
 tcaatgagga cattggcctg gccacagcgt tcaccattgc ccacgagatc gggcacacat  
 1260  
 5 tcggcatgaa ccatgacggc gtgggaaaca gctgtggggc cgtgggtcag gacccagcca  
 1320  
 agctcatggc tgccacatt accatgaaga ccaaccatt cgtgtggtea tctgcagcc  
 1380  
 gtgactacat caccagcttt ctagaactgg gctggggct ctgctgaac aacgggcccc  
 1440  
 ccagacagga ctttgtgtac ccgacagtgg caccgggcca agcctacgat gcagatgagc  
 1500  
 aatgcgcctt tcagcatgga gtcaaatcgc gaggtctgca gcgagctgtg gtgtctgagc  
 1560  
 15 aagagcaacc ggtgcacac caacagcatt ccggcgcgcg agggcacgct gtgccagacg  
 1620  
 cacaccatcg acaaggggtg gtgctacaaa cgggtctgtg tcccttttgg gtgcgcccc  
 1680  
 gaggtgtgtg acggagcctg ggggcctgtg attccaggcg ggactgcagc cggacctgtg  
 1740  
 20 gggggggcgt gtctctctct agcctcact gcgacagcct caggccaacc atcgggggca  
 1800  
 agtctctctt ggtgagaga agggggcacc gctctgcaa caggatgac tgtccccctg  
 1860  
 25 gcccacagga ctccagagaa gtgcagtttt ctgatttga cagatccct ttcgtggga  
 1920  
 attctctaaa gtggaagag tcccaggag cttctctaa attctctg ctgaggtg  
 1980  
 tgggaggtt ctccacatg tctctctaa cttctctaa cttctctaa cttctctaa  
 2040  
 30 ggttctgga cctctctaa attctctaa ggttctgga cctctctaa cttctctaa  
 2100  
 ggttctgga cctctctaa attctctaa ggttctgga cctctctaa cttctctaa  
 2160  
 35 cctctctaa cctctctaa ggttctgga cctctctaa attctctaa cttctctaa  
 2220

cccagcccca ccgtctgctt ctacttgga ccaccttca atgctgacag gggccagacc  
 2400  
 aggtccagag cctcgaagcc ctgggaccga ttaatgcac tctcatcttc atggtgctgg  
 2460  
 5 cccgaccca gctgcctgac ctccgtacc gcttcaatgc cccatctgc cgtgactctc  
 2520  
 tgcctcccta cctctggcac tatggccctt ggaccaagtg ctgcccagt gtgcaggcgg  
 2580  
 tagctgggtg tagggcgttg agtgccgcaa ccaagctgga cagctccggg gtgcgcctcc  
 2640  
 10 actatctgag tgcctacagc aagcttgccc aaaagcaagc ggccttgcaa caggagcct  
 2700  
 tgcctcaaga ctgggttgta ggaactgtg ctctgcagcc gcagcttgcc atgcaaggcg  
 2760  
 15 tgccttgccc ctgggtcttg tgcctaacgc cgcgtctctg ccgcaagaa aaggccttg  
 2820  
 accatgggtg atgcccgcag ccgcccacc ctgtactgag gcttgcacc gccccacttg  
 2880  
 cctctgggag tggcggcctt ccactggtt gactgcacc ccagctggcg gccgggcttc  
 2940  
 20 ccca ccca tggcctttg caagagcgca gaccaccgag ccacgttccc ccgggcgca  
 3000  
 tgcctcccg ccgccaagcc accggccacc atgccttga acttggccc ctccccccc  
 3060  
 25 gcccg tggg tggctggcga ctgggttgag tgcctgcac agtgcggcct cgggcagcgg  
 3120  
 cagctctggg tggctgcac cagccaccg ggcaggggt cgcacagtg caggaggcc  
 3180  
 ctggctccgc ccactagtaa gcttcttga cccgggaatt aattccggac cgttacctgc  
 3240  
 30 aggcgtacca gcttcccta tag  
 3263  
  
 <210> 5  
 35 <211> 1690  
 <212> PRT  
 <213> human

&lt;400&gt; 5

Pro Val Pro Ala Met Pro Gly Gly Pro Ser Pro Arg Ser Pro Ala Pro  
 1 5 10 15  
 Leu Leu Arg Pro Leu Leu Leu Leu Leu Cys Ala Leu Ala Pro Gly Ala  
 5 20 25 30  
 Pro Gly Pro Ala Pro Gly Arg Ala Thr Glu Gly Arg Ala Ala Leu Asp  
 35 40 45  
 Ile Val His Pro Val Arg Val Asp Ala Gly Gly Ser Phe Leu Ser Tyr  
 50 55 60  
 10 Glu Leu Trp Pro Arg Ala Leu Arg Lys Arg Asp Val Ser Val Arg Arg  
 65 70 75 80  
 Asp Ala Pro Ala Phe Tyr Glu Leu Gln Tyr Arg Gly Arg Glu Leu Arg  
 85 90 95  
 Phe Asn Leu Thr Ala Asn Gln His Leu Leu Ala Pro Gly Phe Val Ser  
 15 100 105 110  
 Glu Thr Arg Arg Arg Gly Gly Leu Gly Arg Ala His Ile Arg Ala His  
 115 120 125  
 Thr Pro Ala Cys His Leu Leu Gly Glu Val Gln Asp Pro Glu Leu Glu  
 130 135 140  
 20 Gly Gly Leu Ala Ala Ile Ser Ala Cys Asp Gly Leu Lys Gly Val Phe  
 145 150 155 160  
 Gln Leu Ser Asn Glu Asp Tyr Phe Ile Glu Pro Leu Asp Ser Ala Pro  
 165 170 175  
 Ala Arg Pro Gly His Ala Gln Pro His Val Val Tyr Lys Arg Gln Ala  
 25 180 185 190  
 Pro Glu Arg Leu Ala Gln Arg Gly Asp Ser Ser Ala Pro Ser Thr Cys  
 195 200 205  
 Gly Val Leu Val Tyr Phe Glu Leu Glu Pro Arg Arg Gln Arg Trp Leu  
 210 215 220  
 30 Gln Arg Glu Leu Trp Arg Arg Pro Arg Leu Arg Arg Leu His Gln Ar  
 225 230 235 240  
 Ser Val Ser Lys Glu Lys Trp Val Glu Thr Leu Val Val Ala Asp Ala  
 245 250 255  
 Lys Met Val Glu Tyr His Gly Gln Pro Gln Val Glu Ser Tyr Val Leu  
 30 260 265 270

60

Glu Glu Asp Leu Lys Ile Thr His His Ala Asp Asn Thr Pro Lys Ser  
 305 310 315 320  
 Phe Cys Lys Trp Gln Lys Ser Ile Asn Met Lys Gly Asp Ala His Pro  
 325 330 335  
 5 Leu His His Asp Thr Ala Ile Leu Leu Thr Arg Lys Asp Leu Cys Ala  
 340 345 350  
 Thr Met Asn Arg Pro Cys Glu Thr Leu Gly Leu Ser His Val Ala Gly  
 355 360 365  
 Met Cys Gln Pro His Arg Ser Cys Ser Ile Asn Glu Asp Thr Gly Leu  
 10 370 375 380  
 Pro Leu Ala Phe Thr Val Ala His Glu Leu Gly His Ser Phe Gly Ile  
 385 390 395 400  
 Gln His Asp Gly Ser Gly Asn Asp Cys Glu Pro Val Gly Lys Arg Pro  
 405 410 415  
 15 Phe Ile Met Ser Pro Gln Leu Leu Tyr Asp Ala Ala Pro Leu Thr Trp  
 420 425 430  
 Ser Arg Cys Ser Arg Gln Tyr Ile Thr Arg Phe Leu Asp Arg Gly Trp  
 435 440 445  
 Gly Leu Cys Leu Asp Asp Pro Pro Ala Lys Asp Ile Ile Asp Phe Pro  
 20 450 455 460  
 Ser Val Pro Pro Gly Val Leu Tyr Asp Val Ser His Gln Cys Arg Leu  
 465 470 475 480  
 Gln Tyr Gly Ala Tyr Ser Ala Phe Cys Glu Asp Met Asp Asn Val Cys  
 485 490 495  
 25 His Thr Leu Trp Cys Ser Val Gly Thr Thr Cys His Ser Lys Leu Asp  
 500 505 510  
 Ala Ala Val Asp Gly Thr Arg Cys Gly Glu Asn Lys Trp Cys Leu Ser  
 515 520 525  
 Gly Glu Cys Val Pro Val Gly Phe Arg Pro Glu Ala Val Asp Gly Gly  
 30 530 535 540  
 Trp Ser Gly Trp Ser Ala Trp Ser Ile Cys Ser Arg Ser Cys Gly Met  
 545 550 555 560  
 Gly Val Gln Ser Ala Glu Arg Gln Cys Thr Gln Pro Thr Pro Lys Cys  
 565 570 575  
 35 Lys Gly Arg Tyr Cys Val Gly Glu Arg Lys Arg Phe Arg Leu Cys Asn  
 580 585 590  
 Leu Gln Ala Cys Pro Ala Gly Arg Pro Ser Phe Arg His Val Gln Cys  
 595 600 605  
 Ser His Phe Asp Ala Met Leu Tyr Lys Gly Arg Leu His Thr Trp Val  
 40 610 615 620



61

Pro Val Val Asn Asp Val Asn Pro Cys Glu Leu His Cys Arg Pro Ala  
 625                      630                      635                      640  
 Asn Glu Tyr Phe Ala Glu Lys Leu Arg Asp Ala Val Val Asp Gly Thr  
                          645                      650                      655  
 5 Pro Cys Tyr Gln Val Arg Ala Ser Arg Asp Leu Cys Ile Asn Gly Ile  
                          660                      665                      670  
 Cys Lys Asn Val Gly Cys Asp Phe Glu Ile Asp Ser Gly Ala Met Glu  
                          675                      680                      685  
 Asp Arg Cys Gly Val Cys His Gly Asn Gly Ser Thr Cys His Thr Val  
 10                      690                      695                      700  
 Ser Gly Thr Phe Glu Glu Ala Glu Gly Leu Gly Tyr Val Asp Val Gly  
 705                      710                      715                      720  
 Leu Ile Pro Ala Gly Ala Arg Glu Ile Arg Ile Gln Glu Val Ala Glu  
                          725                      730                      735  
 15 Ala Ala Asn Phe Leu Ala Leu Arg Ser Glu Asp Pro Glu Lys Tyr Phe  
                          740                      745                      750  
 Leu Asn Gly Gly Trp Thr Ile Gln Trp Asn Gly Asp Tyr Gln Val Ala  
                          755                      760                      765  
 Gly Thr Thr Phe Thr Tyr Ala Arg Arg Gly Asn Trp Glu Asn Leu Thr  
 20                      770                      775                      780  
 Ser Pro Gly Pro Thr Lys Glu Pro Val Trp Ile Gln Leu Leu Phe Gln  
 785                      790                      795                      800  
 Glu Ser Asn Pro Gly Val His Tyr Gln Tyr Thr Ile His Arg Glu Ala  
                          805                      810                      815  
 25 Gly Gly His Asp Gln Val Pro Pro Pro Val Ile Ser Trp His Tyr Gly  
                          820                      825                      830  
 Pro Trp Thr Lys Cys Thr Val Thr Tyr Gly Asn Gly Val Gln Arg Gln  
                          835                      840                      845  
 Asn Val Tyr Cys Leu Gln Arg Gln Ala Gly Pro Val Asp Gln Val His  
 30                      850                      855                      860  
 Cys Asp Pro Leu Gly Arg Ile Asp Asp Gln Gln Arg Lys Cys Ser Glu  
 865                      870                      875                      880  
 Gln Pro Cys Pro Ala Arg Trp Trp Ala Gly Glu Trp Gln Leu Cys Ser  
                          885                      890                      895  
 35 Ser Ser Cys Gly Pro Gly Gly Leu Ser Arg Arg Ala Val Leu Cys Ile  
                          900                      905                      910

40

41

42

43

Pro Cys Pro Ala Thr Trp Ala Val Gly Asn Trp Ser Gln Cys Ser Val  
 945 950 955 960  
 Thr Cys Gly Glu Gly Thr Gln Arg Arg Asn Val Leu Cys Thr Asn Asp  
 965 970 975  
 5 Thr Gly Val Pro Cys Asp Glu Ala Gln Gln Pro Ala Ser Glu Val Thr  
 980 985 990  
 Cys Ser Leu Pro Leu Cys Arg Trp Pro Leu Gly Thr Leu Gly Pro Glu  
 995 1000 1005  
 Gly Ser Gly Ser Gly Ser Ser His Glu Leu Phe Asn Glu Ala Asp  
 1010 1015 1020  
 Phe Ile Pro His His Leu Ala Pro Arg Pro Ser Pro Ala Ser Ser Pro  
 1025 1030 1035 1040  
 Lys Pro Gly Thr Met Gly Asn Ala Ile Glu Glu Glu Ala Pro Glu Leu  
 1045 1050 1055  
 15 Asp Leu Pro Gly Pro Val Phe Val Asp Asp Phe Tyr Tyr Asp Tyr Asn  
 1060 1065 1070  
 Phe Ile Asn Phe His Glu Asp Leu Ser Tyr Gly Pro Ser Glu Glu Pro  
 1075 1080 1085  
 Asp Leu Asp Leu Ala Gly Thr Gly Asp Arg Thr Pro Pro Pro His Ser  
 1090 1095 1100  
 20 Arg Pro Ala Ala Pro Ser Thr Gly Ser Pro Val Pro Ala Thr Glu Pro  
 1105 1110 1115 1120  
 Pro Ala Ala Lys Gln Glu Gly Val Leu Gly Pro Trp Ser Pro Ser Pro  
 1125 1130 1135  
 25 Trp Pro Ser Gln Ala Gly Arg Ser Pro Pro Pro Ser Glu Gln Thr  
 1140 1145 1150  
 Pro Gly Asn Pro Leu Ile Asn Phe Leu Pro Glu Glu Asp Thr Pro Ile  
 1155 1160 1165  
 Gly Ala Pro Asp Leu Gly Leu Pro Ser Leu Ser Trp Pro Arg Val Ser  
 1170 1175 1180  
 30 Thr Asp Gly Leu Gln Thr Pro Ala Thr Pro Glu Ser Gln Asn Asp Phe  
 1185 1190 1195 1200  
 Pro Val Gly Lys Asp Ser Gln Ser Gln Leu Pro Pro Pro Trp Arg Asp  
 1205 1210 1215  
 35 Arg Thr Asn Glu Val Phe Lys Asp Asp Glu Glu Pro Lys Gly Arg Gly  
 1220 1225 1230  
 Ala Pro His Leu Pro Pro Arg Pro Ser Ser Thr Leu Pro Pro Leu Ser  
 1235 1240 1245  
 Pro Val Gly Ser Thr His Ser Ser Pro Ser Pro Asp Val Ala Glu Leu  
 1250 1255 1260  
 40

Trp Thr Gly Gly Thr Val Ala Trp Glu Pro Ala Leu Glu Gly Gly Leu  
 1265 1270 1275 1280  
 Gly Pro Val Asp Ser Glu Leu Trp Pro Thr Val Gly Val Ala Ser Leu  
 1285 1290 1295  
 5 Leu Pro Pro Pro Ile Ala Pro Leu Pro Glu Met Lys Val Arg Asp Ser  
 1300 1305 1310  
 Ser Leu Glu Pro Gly Thr Pro Ser Phe Pro Ala Pro Gly Pro Gly Ser  
 1315 1320 1325  
 Trp Asp Leu Gln Thr Val Ala Val Trp Gly Thr Phe Leu Pro Thr Thr  
 10 1330 1335 1340  
 Leu Thr Gly Leu Gly His Met Pro Glu Pro Ala Leu Asn Pro Gly Pro  
 1345 1350 1355 1360  
 Lys Gly Gln Pro Glu Ser Leu Ser Pro Glu Val Pro Leu Ser Ser Arg  
 1365 1370 1375  
 15 Leu Leu Ser Thr Pro Ala Trp Asp Ser Pro Ala Asn Ser His Arg Val  
 1380 1385 1390  
 Pro Glu Thr Gln Pro Leu Ala Pro Ser Leu Ala Glu Ala Gly Pro Pro  
 1395 1400 1405  
 Ala Asp Pro Leu Val Val Arg Asn Ala Ser Trp Gln Ala Gly Asn Trp  
 20 1410 1415 1420  
 Ser Glu Cys Ser Thr Thr Cys Gly Leu Gly Ala Val Trp Arg Pro Val  
 1425 1430 1435 1440  
 Arg Cys Ser Ser Gly Arg Asp Glu Asp Cys Ala Pro Ala Gly Arg Pro  
 1445 1450 1455  
 25 Gln Pro Ala Arg Arg Cys His Leu Arg Pro Cys Ala Thr Trp His Ser  
 1460 1465 1470  
 His Asp Trp Thr Lys Cys Ser Arg Ser Tyr Gly Gly Gly Ser Ser Val  
 1475 1480 1485  
 Arg Asp Thr His Cys Val Asp Thr Arg Asp Leu Arg Pro Thr Arg Trp  
 30 1490 1495 1500  
 Thr His Cys Gln Pro Gly Pro Ala Lys Pro Pro Ala His Arg Pro Cys  
 1505 1510 1515 1520  
 Gly Ala Gln Pro Cys Leu Ser Trp Tyr Thr Ser Ser Trp Arg Glu Cys  
 1525 1530 1535  
 35 Ser Gln Ala Cys Gly Gly Gly Gln Gln Gln Arg Leu Val Thr Cys Pro  
 1540 1545 1550

64

Cys Ser Ala Pro Cys Gly Gly Gly Val Gln Arg Arg Leu Val Lys Cys  
 1585 1590 1595 1600  
 Val Asn Thr Gln Thr Gly Leu Pro Glu Glu Asp Ser Asp Gln Cys Gly  
 1605 1610 1615  
 5 His Glu Ala Trp Pro Glu Ser Ser Arg Pro Cys Gly Thr Glu Asp Cys  
 1620 1625 1630  
 Glu Pro Val Glu Pro Pro Arg Cys Glu Arg Asp Arg Leu Ser Phe Gly  
 1635 1640 1645  
 Phe Cys Glu Thr Leu Arg Leu Leu Gly Arg Cys Gln Leu Pro Thr Ile  
 10 1650 1655 1660  
 Arg Thr Gln Cys Cys Arg Ser Cys Ser Pro Pro Ser His Gly Ala Pro  
 1665 1670 1675 1680  
 Ser Arg Gly His Gln Arg Val Ala Arg Arg  
 1685 1690

15

<210> 6  
 <211> 5338  
 <212> DNA  
 <213> human

20

<400> 6  
 cagggttcctg caatgcccccg cggcccagat ccccgccagcc ccggcgccctt gctggcgccc  
 60  
 ctcttcctgc tcttcggcgc tctggctccc ggcgcccccg gaccgcacc aggaagtgca  
 25 120  
 cccgagggcc cggcgccact ggcacatctg ccccgggttc gactcgacgc cgggggcttc  
 180  
 ctctgtccct acgagctgtg gcccgcgcga ctgcgcaagc gggatgtatc tctggcgcca  
 240  
 30 cccggcgccc cctctacaa cctacaaatc cggggcgccg agctggcctt caacctgacc  
 300  
 cccaatcagc aactgctgcg ccccggcttt gtgagcgaga cggcgggcg cggcgggcctg  
 360  
 ggccggcgcg acatccgggc ctacaccccg gctgcacacc tgcttggcga ggtgcaggac  
 420  
 35 cctgagctcg aggggtggct ggcgccatc agcgctgcg accgctgaa aggtgtgttc  
 480  
 cagctctcca acgaggacta cttcattgag cccctggaca gtgccccggc cgggctcgcc  
 540

caagcccagc cccatgtggt gtacaagcgt caggccccgg agaggctggc acagcggggg  
 600  
 gattccagtg ctccaagcac ctgtggagtg caagtgtacc cagagctgga gcctcgacgg  
 660  
 5 gagegttggg agcagcggca gcagtggcgg cggccacggc tgaggcgtct acaccagcgg  
 720  
 tcggtcagca aagagaagtg ggtggagacc ctggtggtag ctgatgccaa aatggtggag  
 780  
 taccacggac agccgcaggt tgagagctat gtgctgacca tcatgaacat ggtggctggc  
 840  
 10 ctgtttcatg accccagcat tgggaacccc atccacatca ccattgtgcg cctggctctg  
 900  
 ctggaagatg aggaggagga cctaaagatc acgcaccatg cagacaacac cccgaagagc  
 960  
 15 ttctgcaagt ggcagaaaag catcaacatg aagggggatg cccatccctt gcaccatgac  
 1020  
 actgcacatc tctcaccag aaaggacctg tgtgcaacca tgaaccggcc ctgtgagacc  
 1080  
 ctgggactgt cccatgtggc gggaatgtgc cagccgcacc gcagctgcag catcaacgag  
 1140  
 20 gacacggggc tcccgctggc ctccactgta gccacagagc tgggacacag ttttggcatt  
 1200  
 cagcatgacg caagcggcaa tgactgtgag ccggttggga aacgacctt catcatgct  
 1260  
 25 cacagctcc tgcacagcgc cgtccacctc acatgggtc cgtgagagcc ccagtatata  
 1320  
 atcaggttcc ctgacgttg gtggacatca tcttcaaa cctcctgtt caaagacat  
 1380  
 ctccattcc cttccctcc cctcctgtt ctctcgttt tctatcca gttggcctt  
 1440  
 30 agtaccggg ctccacttgc cttctgcag gacatgata atgtctgcca cacactctga  
 1500  
 tgcctgttgg ggaacacctg tcactccaag ctgcatgtag cgttggaagg cacccgctgt  
 1560  
 35 agggagata agtctgtct cagtggggag tggatcccc ttggtttccg gcccgaggcc  
 1620

16 1641

tgtgtgggtg agcgcaagcg ctttcgcctc tgcaacctgc aggcctgccc tgctggccgc  
 1800  
 cctctcttcc gccacgtcca gtgcagccac ttgtagcca tgctctacaa gggccgggtg  
 1860  
 5 caccatggg tgcccggtgt caatgacgtg aacctctggc agctgcactg ccggcccjcg  
 1920  
 aatgagtaact ttgcagagaa gctgggggac gccgtggctg atggcaccac ctgctacag  
 1980  
 gtccagagcca gccgggacct ctgcatcaac ggcattctga agaaccgggg ctgtgacttc  
 2040  
 10 gagattgact ccggtgctat ggaggatcgc tgtgtgtgtg gcacaggcaa cggctccacc  
 2100  
 tgccacaccg tgagcgggac ctttcaggag gccgagggcc tggggatgtt gcatgtgggg  
 2160  
 15 ctgactccag cgggggcacg cgagatccgc atcacaaggc ttgcagagcc tgccaacttc  
 2220  
 ctggcactgc ggagcgagga cccggagaag tacttctcca atgggtgctg gaaccatcag  
 2280  
 tggaaagggg actaccaggt ggcaggacc accttcacat acgcacgcag gggcaactgg  
 2340  
 20 gagaactcca cgtcccggtg tccacccaag gaggctgtct ggatccagct gctgttccag  
 2400  
 gagagcacc ctgggggtga ctacggtac accatccaca gggaggcagg tggccacgac  
 2460  
 25 gagatccgc cggccgtgtt ctcttggcat tatgggccc ttgaccaagt cacagtcacc  
 2520  
 tgggttagag gtgtgcagag gcagaatgtg tactgtcttg agcggcctgg agggccctg  
 2580  
 gaggagagc acgtgaccc cctgggcccg cctgatgacc aacagaggaa gtgcagcgag  
 2640  
 30 cagccttgc ctgccaggtg gttggcaggc gattggcagg tgtgctccag ctctgcggg  
 2700  
 cctgggggcc tctcccgcc ggcgtgtctc tgcacccca gcttgggctt gcatgagag  
 2760  
 35 aggcctcttg agccacccgc ctgtgaacac cttcccccgc cccctactga aacctcttgc  
 2820  
 aaccgcatg taccctgtcc gccacactgg gctgtgggga actggtctca gtgtcagt  
 2880  
 acatgtgggg agggcactca gcgccgaaat gtccctctga ccaatgacac cgggtgtccc  
 2940  
 40

tgtgacgagg cccagcagcc agccagcgaa gtcacctgct ctctgccact ctgttggttg  
 3000  
 cccctgggca cactgggccc tgaaggetca ggcagcggct cctccagcca cgagctcttc  
 3060  
 5 aacgaggetg acttcacccc gcaccacctg gccccacgcc ctccaccgc ctcacaccc  
 3120  
 aagccgggca ccattgggca cgcatttgag gaggaggctc cagagctyga cctgcccggg  
 3180  
 cccgtgtttg tggacgaatt ctactacgac tacaatttca tcaatttcca cgaggatctg  
 10 3240  
 tcctaaggga cctctgagga gcccgatcta gacctggcgg ggacagggga cgggacaccc  
 3300  
 ccaccacaca gccgtcctgc tgcgcctccc acgggtagcc ctgtgcctgc cacagagcct  
 3360  
 15 cctgcagcca aggaggaggg ggtactggga ccttggtccc cgagcccttg gcttagccag  
 3420  
 gccggccgct ccccccccc accctcagag cagacccctg ggaacccctt gatcaatttc  
 3480  
 ctgcctgagg aagacacccc cataggggcu ccagatcttg ggctcccag cctgtccttg  
 20 3540  
 ccagaggttt ccactgatgg cctgcagaca cctgcacccc ctgagagcca aaatgatctc  
 3600  
 ccagtgggca aggacagena cagcagcttc cccctccat ggccgggag gagccaatgag  
 3660  
 25 gtttcaagg atgatgagga aacaaaggac cggggagcac cccacttgc ccgagaccc  
 3720  
 aattatggt tctcccttc atgtctctt ggcacagcc actcctctt tctatctgag  
 3780  
 ctgttgaggt tctggatggt aggaacatc cctacagag caatcttca agtgatctg  
 3840  
 30 aacctgtgaa ccagcgaa tctggtaact gctggggtag ctctctctt tcttctctt  
 3900  
 atgtcccttc tcccagacat gaaggtcagc gatagttccc tggagtcggg caatcccttc  
 3960  
 35 tcccagaccc caggatagg ctataggaa ccagagactg tggcaggtg cagacccctc  
 4020  
 40 4080

ccagcttggg acagccccgc caacagccac agagtccctg agaccacagc gctggctccc  
 4200  
 agcctggctg aagcgggggc ccccgaggac ccgttgjttg tcaggaaagc cagctggcaa  
 4260  
 5 gggggaact ggagcgagtg ctctaccacc tgtggcctgg gtgcggtctg gaggccggtg  
 4320  
 cgctgtagct ccggccggga tgaggactgc gcccccgctg gccggcccca gctgcccgc  
 4380  
 cgctgcacac tggggccctg tgccactgg cactcaggca actggagtaa gtgctcccgc  
 10 4440  
 agctgcggcg gaggttccct agtgccggac gtgcagtgtg tggacacacg ggaacctcgg  
 4500  
 ccactgcggc ccttccattg tcagccggg cctgccaagc cgcctgcgca ccggccctgc  
 4560  
 15 ggggcccagc cctgcctcag ctggtaacaa tcttccctga gggagtgcct ccaggcctgt  
 4620  
 gggggtgggt agcagcagcg tctagtgaac tgcccgagc caggccctctg ccaggaggcg  
 4680  
 ctgaacacca acaccaccgc gccctgcac acccaccct gcacgcagtg gttgggtggg  
 20 4740  
 cccctgggac agtggtcagc cccctgtggt ggtggtgtcc agcggcgccct ggtcaagtgt  
 4800  
 gtcacacacc agacagggtt gcccgaggaa gacagtgaac agtgtggcca ccaggccctg  
 4860  
 25 cctgtgagct cccggccctg tggcaacgac gattgtgagc cgttcagagc cccccgctgt  
 4920  
 gaggcgaac gctgtccct cgggttctga gagaagctgc gcttaactgg cgtctgccag  
 4980  
 ctgcacacca tccgcaccca gtgctgcgcc tctgtctctc ggcacagcca cggcgccccc  
 30 5040  
 tcccagggcc atcagcgggt tgcccgccgc tgactgtgcc aggatgcaca gaccgaccca  
 5100  
 cagacctcag tgcccaccac gggctgtggc ggagctcccg cccctgcgc cctaattggt  
 5160  
 35 ctaacccctt ctcaataccc agcagcaggg tggggaacct ctcccccctc aaaaagggtat  
 5220  
 ttttttatcc taacagtttg tgtaacattt attatgattt tacataaatg agcatctacc  
 5280  
 attcnaaagc aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaa  
 40 5338



69

&lt;210&gt; 7

&lt;211&gt; 947

&lt;212&gt; PRT

5 &lt;213&gt; human

&lt;400&gt; 7

```

Met Gln Phe Val Ser Trp Ala Thr Leu Leu Thr Leu Leu Val Arg Asp
  1             5             10             15
10 Leu Ala Glu Met Gly Ser Pro Asp Ala Ala Ala Val Arg Lys Asp
    20             25             30
Arg Leu His Pro Arg Gln Val Lys Leu Leu Glu Thr Leu Ser Glu Tyr
    35             40             45
Glu Ile Val Ser Pro Ile Arg Val Asn Ala Leu Gly Glu Pro Phe Pro
15     50             55             60
Thr Asn Val His Phe Lys Arg Thr Arg Arg Ser Ile Asn Ser Ala Thr
    65             70             75             80
Asp Pro Trp Pro Ala Phe Ala Ser Ser Ser Ser Ser Ser Thr Ser Ser
    85             90             95
20 Gln Ala His Tyr Arg Leu Ser Ala Phe Gly Gln Gln Phe Leu Phe Asn
    100            105            110
Leu Thr Ala Asn Ala Gly Phe Ile Ala Pro Leu Phe Thr Val Thr Leu
    115            120            125
Leu Gly Thr Pro Gly Val Asn Gln Thr Lys Phe Tyr Ser Gln Gln Gln
25     130            135            140
Ala Gln Leu Lys His Tyr Phe Tyr Tyr Gly Tyr Val Asn Ile Asn Pro
    145            150            155
Gln His Thr Ala Tyr Leu Ser Leu Tyr Ser Ile Met Ile Gly Ile His
    160            165            170
30 Ala Ser His Asp Gly Asp Tyr Phe Ile Gln Ile Leu Gln Ser Met Asp
    175            180            185            190
Gln Gln Glu Asp Glu Glu Glu Gln Asn Lys Pro His Ile Ile Tyr Arg
    195            200            205
Arg Ser Ala Pro Gln Arg Gln Pro Ser Thr Gly Arg His Ala Cys Asp

```

Leu Asn Ser Gly Leu Ala Thr Glu Ala Phe Ser Ala Tyr Gly Asn Lys  
                   260                  265                  270  
 Thr Asp Asn Thr Arg Glu Lys Arg Thr His Arg Arg Thr Lys Arg Phe  
                   275                  280                  285  
 5 Leu Ser Tyr Pro Arg Phe Val Glu Val Leu Val Val Ala Asp Asn Arg  
       290                  295                  300  
 Met Val Ser Tyr His Gly Glu Asn Leu Gln His Tyr Ile Leu Thr Leu  
   305                  310                  315                  320  
 Met Ser Ile Val Ala Ser Ile Tyr Lys Asp Pro Ser Ile Gly Asn Leu  
 10                  325                  330                  335  
 Ile Asn Ile Val Ile Val Asn Leu Ile Val Ile His Asn Glu Gln Asp  
                   340                  345                  350  
 Gly Pro Ser Ile Ser Phe Asn Ala Gln Thr Thr Leu Lys Asn Phe Cys  
                   355                  360                  365  
 15 Gln Trp Gln His Ser Lys Asn Ser Pro Gly Gly Ile His His Asp Thr  
       370                  375                  380  
 Ala Val Leu Leu Thr Arg Gln Asp Ile Cys Arg Ala His Asp Lys Cys  
   385                  390                  395                  400  
 Asp Tor Leu Gly Leu Ala Glu Leu Gly Thr Ile Cys Asp Pro Tyr Arg  
 20                  405                  410                  415  
 Ser Cys Ser Ile Ser Glu Asp Ser Gly Leu Ser Thr Ala Phe Thr Ile  
                   420                  425                  430  
 Ala His Gln Leu Gly His Val Phe Asn Met Pro His Asp Asp Asn Asn  
                   435                  440                  445  
 25 Lys Cys Lys Glu Gln Gly Val Lys Ser Pro Gln His Val Met Ala Pro  
       450                  455                  460  
 Tor Leu Asn Phe Tyr Thr Asn Pro Trp Met Trp Ser Lys Cys Ser Arg  
   465                  470                  475                  480  
 Lys Tyr Ile Thr Glu Phe Leu Asp Thr Gly Tyr Gly Glu Cys Leu Leu  
 30                  485                  490                  495  
 Asn Glu Pro Glu Ser Arg Pro Tyr Pro Leu Pro Val Gln Leu Pro Gly  
                   500                  505                  510  
 Ile Leu Tyr Asn Val Asn Lys Gln Cys Glu Leu Ile Phe Gly Pro Gly  
                   515                  520                  525  
 35 Ser Gln Val Cys Pro Tyr Met Met Gln Cys Arg Arg Leu Trp Cys Asn  
       530                  535                  540  
 Asn Val Asn Gly Val His Lys Gly Cys Arg Thr Gln His Thr Pro Trp  
   545                  550                  555                  560  
 Ala Asp Gly Thr Glu Cys Glu Pro Gly Lys His Cys Lys Tyr Gly Phe  
 40                  565                  570                  575

71

Cys Val Pro Lys Glu Met Asp Val Pro Val Thr Asp Gly Ser Trp Gly  
 580 585 590  
 Ser Trp Ser Pro Phe Gly Thr Cys Ser Arg Thr Cys Gly Gly Gly Ile  
 595 600 605  
 5 Lys Thr Ala Ile Arg Glu Cys Asn Arg Pro Glu Pro Lys Asn Gly Gly  
 610 615 620  
 Lys Tyr Cys Val Gly Arg Arg Met Lys Phe Lys Ser Cys Asn Thr Glu  
 625 630 635 640  
 Pro Cys Leu Lys Gln Lys Arg Asp Phe Arg Asp Glu Gln Cys Ala His  
 10 645 650 655  
 Phe Asp Gly Lys His Phe Asn Ile Asn Gly Leu Leu Pro Asn Val Arg  
 660 665 670  
 Trp Val Pro Gln Tyr Ser Gly Ile Leu Met Lys Asp Arg Cys Lys Leu  
 675 680 685  
 15 Phe Cys Arg Val Ala Gly Asn Thr Ala Tyr Tyr Gln Leu Arg Asp Arg  
 690 695 700  
 Val Ile Asp Gly Thr Pro Cys Gly Gln Asp Thr Asn Asp Ile Cys Val  
 705 710 715 720  
 Gln Gly Leu Cys Arg Gln Ala Gly Cys Asp His Val Leu Asn Ser Lys  
 20 725 730 735  
 Ala Arg Arg Asp Lys Cys Gly Val Cys Gly Gly Asp Asn Ser Ser Cys  
 740 745 750  
 Lys Thr Val Ala Gly Thr Phe Asn Thr Val His Tyr Gly Tyr Asn Thr  
 755 760 765  
 25 Val Val Arg Ile Pro Ala Gly Ala Thr Asn Ile Asp Val Arg Gln His  
 770 775 780  
 Ser Ile Ser Gly Gln Thr Asp Asp Arg Asn Tyr Leu Ala Ile Ser Ser  
 785 790 795  
 Ser Tyr Ala Gln Phe Ile Leu Asn Gly Asn Phe Val Val Thr Ser Ala  
 800 805 810  
 30 Lys Arg Gln Ile Arg Ile Gly Asn Ala Val Val Gln Tyr Ser Gly Ser  
 815 820 825 830  
 Glu Thr Ala Val Glu Arg Ile Asn Ser Thr Asp Arg Ile Gln Gln Glu  
 835 840 845  
 35 Ser Ile Ile Gln Val Leu Ser Val Gly Lys Leu Tyr Asn Pro Asp Val  
 850 855 860

41

72

Glu Arg Lys Arg Lys Leu Val Cys Thr Arg Glu Ser Asp Gln Leu Thr  
                   900                  905                  910  
 Val Ser Asp Gln Arg Cys Asp Arg Leu Pro Gln Pro Gly His Ile Thr  
                   915                  920                  925  
 5 Glu Pro Cys Gly Thr Asp Cys Asp Leu Arg Trp Ala Thr Val Phe Ser  
       930                  935                  940  
 Arg Pro Leu  
 945

10 <210> 8  
     <211> 4086  
     <212> DNA  
     <213> human

15 <400> 8  
     ggtcgttggtg ctggagctta agttgagtag taggaatgcg gtagtagtta ggataatata  
       40  
     tatagttaaa ttaagaatgg ttatgttagg gttgtacggt agaastgcta taattcater  
       80  
 20 tatgttggtta attgagcagp atgctaagat ttgcgctago tgggtatggg taattcater  
       120  
     tcaactggct gctatgctgg ataatatgga gagagtgagg agaaggctta cgtttagtga  
       160  
     aggagagatt tggatatgga ttgagatgga ggtctagttt tgtcatgtga gaagaagcag  
       200  
 25 tccggatgtc agaggggtgc ctgggctaac ctctgggact cagaagtga agggggctat  
       240  
     tcttagtttt attgctatag caattatgat tcttaatgat gactattgat tggtagtatt  
       280  
 30 ggttatgggt caattgctgg agagtatatt gttgaagagg atagctatta gaaggattat  
       320  
     ggatgggggt acttgcgtga ggaataactt gatggcagct tctgttgaac gagggtttat  
       360  
     ttttttgggt agaactggaa taaaagctag catgtttatt tctagacctt ctcaggtaaa  
       400  
 35 aaatcagtcg gagcttagcg ctgtgatgag tgtgcctgca aagatggtag agtagatgac  
       440  
     gggggggggg tgggaagcac catgcagttt gtatcctggg ccacactgct aacgtctctg  
       480

gtgcggggacc tggccgagat ggggagccca gacgcggggg cggccgtggg caaggacagg  
 730  
 ctgcacccga ggcaagtga attattagag accctgagcg aatacgaaat cgtgtctccc  
 840  
 5 atccgagtga acgtctcggg agaacccttt cccacgaacg tccacttcaa aagaacgoga  
 900  
 cggagcatta actctgccac tgacccctgg cctgccttgg cctcctcttc ttcctcctct  
 960  
 acctcctccc aggcgcatta ccgcctctct gccttcgggc agcagtttct attaatctc  
 10 1020  
 accgcatacg ccggatttat cgcctccactg ttcactgtca cctcctcggg gacgcccggg  
 1080  
 gtgaatcaga ccaagtttta ttccgaagag gaagcgggaa tcaagcactg tttctacaaa  
 1140  
 15 ggctatgtca ataccaactc cgagcacacg gcggtcatca gcctctgctc aggaatgctg  
 1200  
 ggcacatccc ggtctcatga tggggattat tttatgaac cactacagtc tatggatgaa  
 1260  
 caagaagatg aagaggaaca aaacaaaccc cacatcattt ataggcgag cgcctccag  
 1320  
 20 agagagcctt caacaggaag gcctgcctgt gacacctcag aacacaaaaa taggcacagt  
 1380  
 aaatctacac agaaaaccag agcaagaaaa tggggacaaa ggaataacct ggctgggtgac  
 1440  
 25 atagagatga taacacaggg cttagcaaca caagcatttt ctgcttatgg taataagacg  
 1500  
 ctatccctat cctcctcagc cctcctcaga atctcctcct cctcctcctc cctcctcctc  
 1560  
 cctcctcctc cctcctcctc cctcctcctc cctcctcctc cctcctcctc cctcctcctc  
 1620  
 30 caaatctcct cctcctcctc cctcctcctc cctcctcctc cctcctcctc cctcctcctc  
 1680  
 ggaactttaa ttaataattt tatgttgaa ctaattgtga ttaataatga ataggatggg  
 1740  
 35 tttctcctat cctcctcctc ttagcacaac ttaaaaaa tttgcctgtg gtaataatga  
 1800  
 40

cccatagaa gctgttctat tagtgaagat agtggattga gtacagcttc tacgatggc  
 1980  
 catgagctgg gccatgtgtt taacatgcct catgatgaca acaacaaatg taaagaagaa  
 2010  
 5 ggagtttaaga gtccccagca tgtcatgget ccaacactga acttctacac caacccctgg  
 2100  
 atgtggtcaa agtgtagtgc aaaatatatc actgagtttt tagacactgg ttatggcgag  
 2130  
 tgttttatta acgaacctga atccagaccc taccctttgc ctgtccaaact gccaggcctc  
 10 2220  
 ctttacaacg tgaataaaca atgtgaattg atttttggac caggtttctca ggtgtgccc  
 2230  
 tatatgatgc agtgcagacg gctctgggtgc aataaagtc atggagtaca caaaggctgc  
 2340  
 15 cggacttaagc acacacccctg ggccgatggg acggagtgcg agcctggaaa gcactgcaag  
 2400  
 tatggatctt gtgttcccaa agaaatggat gtccccgtga cagatggatc ctgggggaagt  
 2430  
 tggagttcct ttggaacctg ctccacaaca tgtggagggg gcacaaaaa agccattcga  
 20 2500  
 gagtgtaaca gaccagaacc aaaaaatggt ggaaaatact gtgtaggacg taqaatgaa  
 2530  
 tttaa tccct gtaacacgga gccatgtctc aagcagaagc gagacttccg agatgaacg  
 2640  
 25 tgtgtctact ttgaggggaa gaattttaac atcaacgggc tgcctcccaa tggcgctgg  
 2700  
 gtcccctaat acagtggaa tctgatgaag gacgggtgca agttgtctct cagagtggca  
 2730  
 gggaaacacg cctactatca gcttcagagc agagtgatag atggaactcc ttgtggccag  
 30 2800  
 gacacacatg atatctgtgt ccagggcctt tgcgggcaag ctggatgca tcatgtttta  
 2830  
 aactcaaaag ccgggagaga taaatgtggg gtttgtgttg gcgataattc ttcattgaaa  
 2940  
 35 acagtggcag gaacatttaa tacagtaaat tatgggtaca atactgtgtt ccgaattcca  
 3000  
 gctggtgcta ccaatattga tgtggggcag cacagttctc caggggaaac agacgatgac  
 3030  
 aactacttag ctttatcaag cagttaaagg gaattcttgc taaatggaaa ctttgtttgc  
 40 3120

acaatggcca aaagggaaat tcgcattggg aatgctgtgg tagagtacag tgggtccgag  
 3180  
 actgccgtag aaagaattaa ctcaacagat cgcattgagc aagaactttt gcttcaggtt  
 3240  
 5 ttgtcgggtgg gaaagttgta caaccccgat gtacgctatt ctttcaatat tccaattgaa  
 3300  
 gataaacctc agcagtttta ctggaacagt catggggccat ggcaagcatg cagtaaaccr  
 3360  
 tgccaagggg aacggaaaacg aaaacttggt tgcaccaggg aatctgatca gcttactggt  
 10 3420  
 ttctgatcaa gatgcgacag gctgcccag cctggacaca ttactgaacc ctgtgggtaca  
 3480  
 gactgtgacc tgaggtyggc cactgttttc tcaaggcctt tataaatgaa ttgtgagagt  
 3540  
 15 cttgcaggag gcccagcag gagaagcaaa aggaggygat gccggtcttt agttccctt  
 3600  
 tcttggtttt cagtgaata agctttaacc aattctccat cctctggaa ctgattatcc  
 3660  
 aagacataca tttgcagatt tcttggtcac ctaagaanta aaaatagcta atagagtatg  
 20 3720  
 gcaattgcca aaaaaaatc agttgatccr caacacttcc tgggtaggta ttageattat  
 3780  
 tcttgagttc catgttaagt gaaaacttgt ttgaaaactr aaagaaaag agggagaacc  
 3840  
 25 tttatcttca agttatccat aatgactat atctacccaa agttatccr aaccagtaca  
 3900  
 tttatcttca agttatccat aatgactat atctacccaa agttatccr aaccagtaca  
 3960  
 30 tttatcttca agttatccat aatgactat atctacccaa agttatccr aaccagtaca  
 4020  
 tttatcttca agttatccat aatgactat atctacccaa agttatccr aaccagtaca  
 4080  
 35 tttatcttca agttatccat aatgactat atctacccaa agttatccr aaccagtaca  
 4140

<220>

<223> primer

5 <400> 9

gatcgcgggcc gctatgggtgg acacgtggcc tctatggctc c

41

<210> 10

10 <211> 43

<212> DNA

<213> Artificial Sequence

<220>

15 <223> primer

<400> 10

tgaggcccttc agggccgata actgtgcaga gcactcacc cct

43

20



Claims

1. An MPTS protein selected from the group consisting of MPTS-15, MPTS-10,  
MPTS-19 and MPTS-20, wherein said protein is present in other than its natural  
5 environment.
2. The protein according to claim 1, wherein said protein has an amino acid  
sequence substantially identical to the sequence of SEQ ID NO:01, 03, 05 or 07.
- 10 3. A nucleic acid present in other than its natural environment, wherein said  
nucleic acid has a nucleotide sequence encoding an MPTS protein selected from the  
group consisting of MPTS-15, MPTS-10, MPTS-19 and MPTS-20.
4. A nucleic acid according to claim 3, wherein said nucleic acid has a nucleic acid  
15 sequence that is the same as or substantially identical to the nucleotide sequence of SEQ  
ID NO:02, 04, 06 or 08.
5. An expression cassette comprising a transcriptional initiation region functional  
in an expression host, a nucleotide sequence according to claims 3 or 4 under the  
20 transcriptional regulation of said transcriptional initiation region, and a transcriptional  
termination region functional in said expression host.
6. A cell comprising an expression cassette according to claim 5 as part of an  
extrachromosomal element or integrated into the genome of a host cell as a result of  
25 introduction of said expression cassette into said host cell.
7. The cellular progeny of the host cell according to claim 6.
8. A monoclonal antibody binding specifically to an MPTS protein according to  
30 claim 1.

isolating said protein substantially free of other proteins.

10. An MPTS protein as claimed in claim 1 or 2, whenever produced by the process of claim 9.

5

11. A method of screening to identify MPTS modulatory agents, said method comprising:

contacting an MPTS protein according to claim 1 with a substrate in the presence of an potential modulatory agent; and

10 determining the effect of said modulatory agent on the activity of said protein.

12. The method according to claim 11, wherein said substrate comprises a glu-ala bond.

15 13. The method according to claim 12, wherein said substrate is aggrecan or a fragment thereof.

14. A method of treating a host suffering from a disease condition associated with MPTS activity specifically wherein said disease condition is characterized by the presence of aggrecan cleavage products, said method comprising:

20 administering to said host an MPTS modulatory agent, specifically an antagonist.

15 15. Use of a MPTS modulatory agent, obtainable or obtained by the method claimed in claim 11 for the preparation of a medicament for the treatment of a disease condition associated with MPTS activity, specifically wherein said disease condition is characterized by the presence of aggrecan cleavage products, like arthritis.

[illegible]

GTCTCTGCTGGTGCTTCTCCCAAGACTATCTTGAAGGTGGGCTGTTTGCCTTTCGTGAACA  
CATTCTTGGTAT  
(SEQ ID NO:02)

FIG. 1B

&gt;ORF(frame +1)

```
5 MEILWKTLTWILSLIMASSEFHS DHRLSYSSQEEFLTYLEHYQLTIPIRVDQNGAFLSFTV
  KNDKHSPPRRRSM DPIDPQQAVSKLFFKLSAYGKHFHLNLTNTDFVSKHFTVEYWGKDGFPQ
  WKHDFLBNCHYTGYLQDQFSTTKVALSNCVGLHGVIATEDEEYFIEPLENTTEDSKHFSYE
  NCHPHVIYKKSALQQEHLVDHSECGVSDFTESGKPWWLNDTSTVSYSLPINNTHIHRQKE
  SVSIERFVETLVVADEMMVGYHGRKDIEHYILSMNIVAKLYRDSSLGNVNIIVARLIVL
10 TEDQPMIEINHHADKSLDSFCKWQKSLSHQSDGNTIPENGIAHHDNAVLTTRYDICTYKN
  KPGDTLGLASVAGMCEPERSCSINEDIGLGSAPTIAHEIGHNFGMNHDCIGNSCGTGHEA
  AKLMAAHITANTNPFSWSACSRYITSFLDSGGGTOLDNEPPKRDFLYFAVAPGQVYDADE
  QCRFQYGATSRQCKYGEVCRELWCLSKSNRCVTNSIPAAEGTLQQTGNIEKGWCYQGDCVF
  FGTWPQSIDGGWGPWSLWGECSFTGGGVSSSLRHCDSPAPSGGGKYCLGERKRYRSCNTD
15 PCPLGSFDFREKQCADFDNMPFEGKYYNWKPYTGGGVKPCALNCLAEGYNFYTERAPAVID
  GTQCNALSLDICECKHVSCDNILGSDAREDRCRVCGGDGSTCDALGFFNDLSLPRGGY
  MEVWQIPRGSVHIEVREVAMSKNYIALKSEGDDYYINGAWTIDWPRKFEVAGTAFHYKRPT
  DEPESEALGPTSENLI VMVLLQEQLNGIRYKFNVPITRTGSGDNEVGFTWNHQSWSSECSA
  TCAGGKMPTRQPTQRRARWRTKHILSYALCLLKKLIGNISCRFASSCNLEKETLL*LYYIPF
20 VFNL M*FVQICW*NTSWHNSCLCNCFSQDYLEGGGLFAFREHILG
  (SEQ ID NO (1))
```

FIG. 1C

## Align MP15 with ADAMTS-6(in public data base)

```

5  MP15-4universal+1_ORF1 MEILWKTLTWILSLIMASSEFHSOHRLSYSSQEEFLTYLEHYQLTIPIRV
   ADAMTS6+1_ORF1        MEILWKTLTWILSLIMASSEFHSOHRLSYSSQEEFLTYLEHYQLTIPIRV
                           *****

10 MP15-4universal+1_ORF1 DQNGAFLSFTVKNDKHSRRRRSMDPIDPQAVSKLFFFLJAYGKHFLNL
   ADAMTS6+1_ORF1        DQNGAFLSFTVKNDKHSRRRRSMDPIDPQAVSKLFFFLJAYGKHFLNL
                           *****

15 MP15-4universal+1_ORF1 TLNTDFVSKHFTVEYWGKDGQPQWKHDFLDNCHYTGVLQDQJSTTKVALSN
   ADAMTS6+1_ORF1        TLNTDFVSKHFTVEYWGKDGQPQWKHDFLDNCHYTGVLQDQJSTTKVALSN
                           *****

20 MP15-4universal+1_ORF1 CVGLHGVATIEDDEEYFIEPLEKNTTEDSKHFSYENGHPHVIVKKSALQQRH
   ADAMTS6+1_ORF1        CVGLHGVATIEDDEEYFIEPLEKNTTEDSKHFSYENGHPHVIVKKSALQQRH
                           *****

25 MP15-4universal+1_ORF1 LYDHSHCGVSDFTSGKPPWLNDDTSTVSYALPINNTHIHHAQKESVSIER
   ADAMTS6+1_ORF1        LYDHSHCGVSDFTSGKPPWLNDDTSTVSYALPINNTHIHHAQKESVSIER
                           *****

30 MP15-4universal+1_ORF1 FVETLWVADPMMVGYHGRKDIHYILSVNNIVAKLYRISALGNVNIIVA
   ADAMTS6+1_ORF1        FVETLWVADPMMVGYHGRKDIHYILSVNNIVAKLYRISALGNVNIIVA
                           *****

35 MP15-4universal+1_ORF1 DNNAVLTTRYICITYFNKPCSTGLGLASVAGHCEPERSCSINEDIOLGSAFT
   ADAMTS6+1_ORF1        DNNAVLTTRYICITYFNKPCSTGLGLASVAGHCEPERSCSINEDIOLGSAFT
                           *****

40 MP15-4universal+1_ORF1 LAHEIGHNPKNNHDOGTGNSDGLRHPAAPHKQAHIAHNTAFSALNFF
   ADAMTS6+1_ORF1        LAHEIGHNPKNNHDOGTGNSDGLRHPAAPHKQAHIAHNTAFSALNFF
                           *****

45 MP15-4universal+1_ORF1 FVETLWVADPMMVGYHGRKDIHYILSVNNIVAKLYRISALGNVNIIVA
   ADAMTS6+1_ORF1        FVETLWVADPMMVGYHGRKDIHYILSVNNIVAKLYRISALGNVNIIVA
                           *****

50 MP15-4universal+1_ORF1 GTWPQSIDGLNGPWSLWGECSFTCGGGVSSSLRHCI SPAISGGGKYCLGE
   ADAMTS6+1_ORF1        GTWPQSIDGLNGPWSLWGECSFTCGGGVSSSLRHCI SPAISGGGKYCLGE
                           *****

55 MP15-4universal+1_ORF1 EKRYFSCNTLPCPLGSRDFREKQCADFNNIPFRGKYYNWFYTGGGVVKPC
   ADAMTS6+1_ORF1        EKRYFSCNTLPCPLGSRDFREKQCADFNNIPFRGKYYNWFYTGGGVVKPC
                           *****

```

FIG. 1C (Cont)

Align MP15 with ADAMTS-6(in public data base):

```

5  MP15-4universal+1_ORF1 ALNCLAFGYNFYIERAPAVIDGTQCNADSLDICECKHVGCNINILGSD
   ADAMTS6+1_ORF1        ALNCLAFGYNFYIERAPAVIDGTQCNADSLDICECKHVGCNINILGSD
                               *****

10 MP15-4universal+1_ORF1 AREDRRCVCGGDSSTCDAIEGFFNDSLPRGGYMEVVQIPRGSVHIEVREV
   ADAMTS6+1_ORF1        AREDRRCVCGGDSSTCDAIEGFFNDSLPRGGYMEVVQIPRGSVHIEVREV
                               *****

15 MP15-4universal+1_ORF1 AMSKNYIALKSEGDDYYINGAWTIDWPRKFDVAGTAFHYKRPTDEPESLE
   ADAMTS6+1_ORF1        AMSKNYIALKSEGDDYYINGAWTIDWPRKFDVAGTAFHYKRPTDEPESLE
                               *****

20 MP15-4universal+1_ORF1 ALGPTSENLIWM/LLQEQLGIRYKFNVPITRTGSGINEVGFTWNHQSW
   ADAMTS6+1_ORF1        ALGPTSENLIWM/LLQEQLGIRYKFNVPITRTGSGINEVGFTWNHQSW
                               *****

25 MP15-4universal+1_ORF1 NLPKETLL (SEQ ID NO:01)
   ADAMTS6+1_ORF1        NLAKEFLL (GenBank Accession No. AF140674)
                               *****

```





TTGCGATGCAAGGCGTGGGCAGCCGCTCGGTCTGTGCGAAGCGCCGCGTCTCTGCGCGA  
AGAAAAAGCGCTGGAACGACAGCGCATGCCCGCAGCCGCGCCACCTGTACTGAGCGCTGCC  
ACGCGCCCACTTGCCCTCCGGAGTGGCGGCGCTCGACTGGTCTGAGTGCACCCCGAGCTGC  
GGGCGGGCGCTCCGCGCACCGCGTGGTCCCTTTGCAAGAGCGCAGACCACCGCGCGCAAGCTGC  
5 CCGCGGGCGCACTGCTCACCCGCGCGCAAGCGACCGGCCACCATGCGCTGCAACFTGGCGCG  
CTGCCCCCGGGCCCGCTGGGTGGCTGGCGAGTGGGGTGAGTGTCTGACACAGTGGGGCGTC  
GGGCGAGCGGCAGCGCTCGGTGCGCTGCACCGCCACACGGGCCAGGCGTCGCAAGAGTGCA  
CGGAGGCGCTGCGGCGCGCGACTAGTAAGCTTCGTCGACCCGGGAATTAATTCCGGACCGG  
TACCTGCAGGCGTACCAGCTTCCCTATAG  
10 (SEQ ID NO: 04)

FIG. 2B

>MP10-full-length+3\_ORF1 Translation of MP10-full-length in frame +3, ORF 1,  
threshold 50

```

5  MAPAQQILRWALALGIGMFEVTHAFRSQDEFLSSLESYEIAFPTEVDHNGALLAFSPPPP
  ERQRRGTGATAESFLFYKVASPSTHFLNLTRSSFLLAGHVSVEYWTREGLAWQRAARPHC
  LYAGHLQGQASSSHVAISTCGGLHGLIVADEEEYLIPLHGGPKGSRSPPEESGPHVVYKRS
  SLEHPHLDTACGVFDEKPKWGRPWWRRLTKPPPAFPLGNETERGQPGPKRSVSRERYVETL
10  UVADKMTVAYHGERDVEQYVLAIMNIVAKLFQDSSLGSTVNILVTFLLILLTEQPTLEITH
  HAGKSLDSFCKWQKSIYNHSGH3NAIPENGVANHDTAVLITRYDICIYKNKPCGTGLAPV
  GGMCEERERSCSVNEDIGLATAFTIAHEIGHTFGMNHDGVGNSCCAFGQDPAKLMAAHITMK
  TNFFVWSSCSRDIYTSFLDSGLGLCLNNRPPRQDFVYPTVAPGQAYDADEQCRFQHGKSR
  GLQRAVVSEQEQPVHHQQHPGGRGHAVPDAHHRQGVVLQTGLCPWVAPRGCGFSLGAVDS
15  MGI CSRTCGGGVSSSSSEHCDSPRPTIGGHYCLGEERRHRSCNTDDCPPGSQDFREVCSEF
  DSIIFRGKFYKWTYRGGGVKACSLTCLAEGFNFYTERAAAVVDGTPCFPDTVDICVSGEC
  KHVGCDRVLGSDLFEDKCRVCCGDSACHTIEGVFSPASPGAGYELVWVLPKGSVHIFIQD
  LNLSLSHLALKGDQESLLEGLPGTPQPHFLPLAGTTFQLRQGPDCVQSLEALGPINASLI
  VMVLARTELPALFYRENAPIAFDLSLPYSVHYAPWTKCSPSVQAVARCERWSAATKLDSSA
20  VAFHYCSAHSKLAQKQARIQHGAIPQDWVVGTVLALQPQLAMQGVRSRSVVCQAPRLCREEK
  ALDDSAQPQPRPEVLRPATAPLALRSGGPRIV*

```

(SEQ ID NO:03)

25

MP19

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40

[illegible]

For the purpose of this paper, we will

FIG 3B

>MP19-full-length+1\_ORF1 Translation of MP19-full-length in frame +1, ORF 1,  
threshold 50

```

5  PVPAMPGGPSPRSPAPLLRPLLLLLCALAPGAPGAPGRATEGRAALDIVHPVRVDAGGSF
   LSYELWPRLARKRDVSVRRDAPAFYELQYRGRELRFNLTAHQHLLAPGFVSETRRRGGLGP
   AHIRAHTPACHLIGEVQDPELEGGLAAISACDGLKGVFQLSNEDYFIEPLDSAPARPGHAQ
   PHVVKPQAPERLAQRGDSSAPSTCGVQVYPELEPRRERWEQRQQWRRPRLRLHQRSVSK
10  EKWVEFLVVADAFMVEYHGQPQVESYVLTIMNMVAGLFHDPSIGNPIHITIVRLVLEDEE
   EDLKHITHADNTPKSFCKWQKSINMKGDAHFLHHDTAILLTKDLCCATMTRPCETLGLSHV
   AGMCQPHRSCSINEDTGLPLAFTVAHELGHSEFGIQHDGSGNDCEPVGKRPFIMSPQLLYDA
   APLTWSECSRQYITRFLDRGWGLCLDDPPAKDIIDFSPVPPGVLYDVSHQCRLQYGAYSAF
   CEDMDHVCHTLWCSVGTTCHSKLLAAVDGTRCGENKWCLSGECVFVGFRPEAVDGGWGSWS
15  AWSICSFSCGMGVQSAERQCTQPTPKYKGRYCVGERKRFRLLCNLQACAPAGRPSFRHVQCCH
   FQAMLYEGRLHTWVPVVNDVNPCELHCFPAKEYFAEKLEDAVVDGTFCYQVRASRDLCING
   ICKT/AGCDFEIDSGAMEDFCGVCHGNGSTCHTVSGTFEEAEGLGYYVDVGLIPAGAREIRIQ
   EVAAANFLALRSEDPEKYFLNGSWTIQWNGDYQVAGTTFTYARRGNWENLTSFGPTFEPV
   WIQILFQESNPGVHYEYTIHREAGGHCEVFPVFSWHEYGEWTKCTVTCGRGVQRQNVYCLE
20  RQAGPVDEEHCDPLGRPDQQRKCEQPCPARWWAGEWQLCSSSCGPGGLSRRAVLICIRSV
   GIDEQSALEPPACEHLPRPPTETPCNHFVPDPATWAVGNWSQCEVTCGEGTQCFEN/LCTND
   TGVPCDEAQQPASEVTCSLPLCRWPLDGTGLGPEGSGSGSSSHELFEADFI PHHLAPRPSPA
   SSKPKPGTMMNAIEEEAPELDLPBPVTVDDFYDYNFINFHEILSYGPSEEPDLDLAGTGDR
   TFPPHSPPAAPSTGSPVPATEPPAAKEEGVLGPWSPSPWFSQAGRSPPPSEQTTCGNFLIN
25  FLPHEDTPIGAPDLGLPLSLWPRVSTDCLOTPATPESQNDFFVGKDSQSQLPPPWRDETNH
   VFKDDEEPKGRGAPHLPPRPSSTLPLLSFVISTHSSSPDVAELMTSGTVAWEPALBGLG
   PVDSELVPTVGVASLLPPPAPLPMEKVEDSSLEPGTSPFPAPGPGSWDLQTVAVWGTFPLP
   TTLTGLCHMPEPALNPGPKSQPESLSPEVPLSSRIISTFANDSPANSHRVPETQPLAPSLA
   EAGPPAPPLVVRNASWQAGNASECSTICGLGAVWRPVRCSSGRDEDAPAGRFPQPARACHL
30  RPCATWHSNWSKCSRSCGSSSVRDVQCVDTRDLEPLLPFHCQPGPAKPPAHRPQGAQPC
   LSWYTSFWRECSEACGGGEQRLVTCEPGLCREALRPNTTEPCNTHPCTQWVVGPGWQCS
   APGCGGVQRRLVKECVNTQTGLPEEDSTIQGHEAWPESSEPCGTEDCEPVEPPRCERDLSE
   AFCETELELIGRCQLPTLFTQCRSCSPPSHCAPSRGHQVAPP*

```

55 SEQ ID NO:051

MP-20

[illegible]

CGATCATGTTTTAAACTCAAAAAGCCCGGAGAGATAAATGTGGGGTTTGTGGTGGCGATAAT  
TCTTCATGCAAAAACAGTGGCAGGAACATTTAATACAGTACATTATGGTTACAATACTGTGG  
TCCGAATTCCAGCTGGTGCTACCAATATTGATGTGCGGCAGCACAGTTTCTCAGGGGAAAC  
AGACGATGACAACACTACTTAGCTTTATCAAGCAGTAAAGGTGAATTCTTGCTAAATGGAAAC  
5 TTTGTTGTCCACAATGGCCAAAAGGGAAATTCCGATTGGGAATGCTGTGGTAGAGTACAGTG  
GCTCCGAGACTGCCGTAGAAAAGAATTAACCAACAGATCGCATTGAGCAAGAACTTTTGCT  
TCAGGTTTGTGCGGTGGGAAAGTTGTACAACCCCGATGTACGCTATTCTTTCAATATTCCA  
ATTGAAGATAAAACCTCAGCAGTTTCTACTGGAACAGTCATGGGCCCATGGCAAGCATGCAGTA  
AACCCTGCCAAGGGGAACGGGAAACGAAAACCTGTTTGCAACAGGGAAATCTGATCAGCTTAC  
10 TGTTTCTGATCAAAGATGCGATCGGCTGCCCCAGCCTGGACAATTACTGAACCTGTGGT  
ACAGACTGTGACCTGAGGTGGGCCACTGTCTTTCTCAAGGCCCTTTATAAATGAATTGTGAGA  
GCTTGGCAGGAGGTCCCAACAGGAGAAGCAAAAGGAGGGGATGCCGGTCTTTAGTTCCCT  
TTCTTGTGTCTTCAGTGAATAAGCTTTAAACCAATCTCTCATCCCTCTGGAACTGATTATCC  
AAGACATACATGTGCAGATTCTTGTTCACCTAAGAAATAAAAATAGCTAATAAGTATGG  
15 CACTTGGCAAAAAAATTCAGTTGATCCTCACMACTTCTGTGGTAGGTATTAGCATTATGA  
TTGAGTCAATTTGTACGTGAAAACTTGTTTGGAAAGTCAAAAGAAAAAGAGGGAGAACCTCA  
TCCTCAAAAGTACCCATAATGACCTATATCTACCGAGAGTGTATACCAACCAGTAGAAGAA  
CTCCTACACACCTGAAAGTTGCAGTACACTAAAGTAGCTCATGGAGGAAACAAGAAGAAA  
ATGTATATATGGATGTYTGAATATTCAACAATTCTGTGTTTAAAGAAAAAAAAAAAAAA  
20 AAAAAAAAAAAAAAAGTACTCTGCGTTGTTACCACTGCTTGCCCTATAGTGAAGTGTATT  
(SEQ ID NO:08)

FIG. 4B

>MP20-full-length-1\_ORF1 Translation of MP20-full-length in frame +1, ORF 1, threshold 150

```

5  MQFVSWATLLTLLVRDLAEMGSPFAAAAVEKDRLHPRQVKLLETLSYEYEVSPIRVNALGE
  PFPTNVHFKRTRRSINSATDPWPAFASSSSSSTSSQAHYFLSAFGQQFLFNLTAINAGFIAP
  LFTVTLLGTPEGVNQTKFYSEEEAELKHCFYKGYVNTNSEHTAVISLCSGMLGTFFESHG DY
  FTEPLQSMDEQEDEEEQNKPHTIYPRSA PQREPSTGRHACDTSEHKNRHSKDKKXTRARKW
10 GERINLAGDVAAALNSGLATEAFSA YGNKTDNTEKRTHRTKRFLSYPRFVEVLVADNEM
  VSYHGENLQHYILTLMSIVASIYHDP SIGNLINIVIVNLIVIHNEQDGPSISFNAQTTLKN
  PQQWQHSKNSPGGIHHDTAVLLTHQD ICRAHDKCDTLGLAELGTICDPYRSCSISGDSGLS
  TAFTIAHELGHVFNMPHDDNNKCHEEGV KSPQHVMAPTLNFTNPMWWSKSRKXITEFLD
  TGYGEULLNEPESRPYPPLPVQLPGI LYNVIMQCELIFGPGSQVC PYMMQCRRLWNNVNGV
15 HKGCRTQHTPWADGTECEPGHCHYGF CVPHEMDVPVTDGSGWSWSPFGTCSRTOGGGIKT
  AIRECNEPEPKNGGKYCVGREMKFHS CNTEPCLKQKRDFRDEQCAHFDGFFHFNINGLLPNV
  RWVPQYSGILMKDRCKLFCRVAGNTA YYQLRDEVIDGTPEGQDTNDICVQGLCRQAGCDHV
  LNSKARREKCGVCGDNSSCKTVAGT FNTVHYGYNTVVRIPAGATNIDVEQHSFSGETDDD
  NYLALSSSEGEPLNGNFVVTMAHREI RIGNAVEYSGETAVERINSTDRIEQELLQVL
20 SVGKLYNPVRYSFNIPIDEPQQFYWN SHGPWQACSKPCQGERKRLVCTRESQQLTVSD
  QRCDALPQPGHITEPCGTDCDLRWAT VFSRPL*
  (SEQ ID NO:07)

```